| Prepared by:   | Dillon Alegre, Grant Brink & Rachel Ellison, Environmental Assessment Services, LLC        |
|----------------|--|
| Report Period: | May 1 <sup>st</sup> to May 15 <sup>th</sup> , 2023   |
| Re:            | CRAMER FISH SCIENCES - WILLAMETTE VALLEY FISH PASSAGE<br>MONITORING VIA ROTARY SCREW TRAPS |

# Project Schedule

| Site  | Task             | Start     | End        | Days |  |  |  |  |  |
|---|------------------|-----------|------------|------|--|--|--|--|--|
| Breitenbush River RST                                   | Trap Install     | TBD       | TBD        | 1    |  |  |  |  |  |
| Breitenbush River RST                                   | Operation        | TBD       | 11/30/2023 | TBD  |  |  |  |  |  |
| Detroit Head of Reservoir- North Santiam River RST      | Trap Install     | 4/19/2023 | 4/19/2023  | 1    |  |  |  |  |  |
| Detroit Head of Reservoir- North Santiam River RST      | Operation        | 5/4/2023  | 11/30/2023 | 210  |  |  |  |  |  |
| Green Peter Head of Reservoir- Middle Santiam River RST | Highline Install | 4/25/2023 | 4/25/2023  | 1    |  |  |  |  |  |
| Green Peter Head of Reservoir- Middle Santiam River RST | Trap Install     | 4/26/2023 | 4/26/2023  | 1    |  |  |  |  |  |
| Green Peter Head of Reservoir- Middle Santiam River RST | Operation        | 5/4/2023  | 11/30/2023 | 210  |  |  |  |  |  |
| Hills Creek Head of Reservoir RST                       | Trap Install     | 5/9/2023  | 5/9/2023   | 1    |  |  |  |  |  |
| Hills Creek Head of Reservoir RST                       | Operation        | 5/9/2023  | 6/30/2023  | 52   |  |  |  |  |  |

#### Table 1. Project Schedule

# Table 2. Sampling Dates for Reporting Period

| Site   | Total<br>Sampling<br>Period Start | Current<br>Reporting<br>Period Start | Current<br>Reporting<br>Period End | Days<br>Sampled This<br>Period | Total<br>Days<br>Sampled |
|--|-----------------------------------|--------------------------------------|------------------------------------|--------------------------------|--------------------------|
| Detroit Head of Reservoir- North<br>Santiam River RST      | 5/4/2023                          | 5/1/2023                             | 5/15/2023                          | 11                             | 11                       |
| Green Peter Head of Reservoir-<br>Middle Santiam River RST | 5/4/2023                          | 5/1/2023                             | 5/15/2023                          | 11                             | 11                       |
| Hills Creek Head of Reservoir RST                          | 5/9/2023                          | 5/1/2023                             | 5/15/2023                          | 6                              | 6                        |

|  |         | =                              |                                     |                |                     |
|--|---------|--------------------------------|-------------------------------------|----------------|---------------------|
| Site   | Species | Catch<br>(Reporting<br>Period) | Recaptures<br>(Reporting<br>Period) | Total<br>Catch | Total<br>Recaptures |
| Detroit Head of Reservoir- North Santiam<br>River RST      | CHS     | 2,098                          | 0                                   | 2,098          | 0                   |
| Detroit Head of Reservoir- North Santiam<br>River RST      | STW     | 114                            | 0                                   | 114            | 0                   |
| Green Peter Head of Reservoir- Middle<br>Santiam River RST | CHS     | 20                             | 0                                   | 20             | 0                   |
| Green Peter Head of Reservoir- Middle<br>Santiam River RST | STW     | 1                              | 0                                   | 1              | 0                   |
| Hills Creek Head of Reservoir RST                          | CHS     | 25                             | 0                                   | 25             | 0                   |

#### Table 3. Willamette Valley Rotary Screw Trap Monitoring Catch Summary

# Summary of Rotary Screw Trap Data

There are 3 rotary screw traps (RSTs) that have been installed and commenced fishing during the reporting period. For this reporting period, traps were operated at the following 3 locations: Detroit Head of Reservoir – North Santiam River, Green Peter Head of Reservoir – Middle Santiam River and Hills Creek Head of Reservoir on the upper Middle Fork Willamette River.

The Detroit Head of Reservoir – North Santiam RST and Green Peter Head of Reservoir – Middle Santiam RST were installed on April 19<sup>th</sup> and 26<sup>th</sup>, respectively. The RSTs at Detroit Head of Reservoir – North Santiam and Green Peter Head of Reservoir – Middle Santiam rivers started sampling on May 4<sup>th</sup> once permits were received. The Hills Creek Head of Reservoir RST on the upper Middle Fork Willamette River was installed and began sampling on May 9<sup>th</sup>.

Winter Steelhead may be present at the Breitenbush River, Detroit Head of Reservoir – North Santiam, and Green Peter Head of Reservoir – Middle Santiam River sites. All natural origin juvenile O. mykiss captured at these sites will be treated and reported as Winter Steelhead.

The RST for the Breitenbush River sampling site is still being constructed by EG Solutions. This RST and its highline are slated for install as soon as the new RST is available.

Sampling start dates are included in Table 2, and season total collection numbers are displayed in Table 3. The locations of the RSTs are depicted in Figures 1 through 4.



Portland Salem Eugene OREGON FIGURE 1 Breitenbush River

**RST** Locations

•

500 Feet








FIGURE 2 North Santiam Above Detroit



•

N





FIGURE 3 Middle Santiam River



•

Imagery Source: 2019, ESRI.



**FIGURE 4** Middle Fork Willamette Above Hills Creek

NFS-Rd21

**Middle Fork** 

Willamette Above **Hills Creek RST** 



NFS-Rd-2120

**RST** Locations

•

**500 Feet** 

He Fork Willamette River



# **Breitenbush River**

The Breitenbush River RST is still being manufactured by EG Solutions. The trap is slated for install as soon as it is available.

#### **Target Species**

No target species have been captured as the RST has yet to be installed. There were 0 Chinook and 0 Winter steelhead caught during the reporting period (Figure 5). Figure 6 shows length frequency data todate. Table 4 provides life stage, length, and weight data for all Chinook Salmon and Winter Steelhead that have been caught at the Breitenbush River site to-date and for the reporting period.




Figure 5. Chinook Captured per day 5/01/2023 to 5/15/2023 (Breitenbush River)




Figure 6. Length Frequency of Juvenile Chinook Sampled Season To-Date (Breitenbush River)

|                      | To-Date |         |       |           |     |          |      |     |                         |      |  |
|----------------------|---------|---------|-------|-----------|-----|----------|------|-----|-------------------------|------|--|
| Site                 | Route   | Species | Life  | Collected | L   | ength (m | m)*  |     | Weight (g) <sup>*</sup> |      |  |
|                      |         |         | stage |           | Min | Мах      | Mean | Min | Max                     | Mean |  |
|                      |         | CHS     | Fry   | 0         | N/A | N/A      | N/A  | N/A | N/A                     | N/A  |  |
|                      |         | CHS     | Parr  | 0         | N/A | N/A      | N/A  | N/A | N/A                     | N/A  |  |
| Breitenbush<br>River | 5ft     | CHS     | Smolt | 0         | N/A | N/A      | N/A  | N/A | N/A                     | N/A  |  |
|                      |         | STW     | Fry   | 0         | N/A | N/A      | N/A  | N/A | N/A                     | N/A  |  |
|                      |         | STW     | Parr  | 0         | N/A | N/A      | N/A  | N/A | N/A                     | N/A  |  |
|                      |         | STW     | Smolt | 0         | N/A | N/A      | N/A  | N/A | N/A                     | N/A  |  |

#### Table 4. Descriptive Statistics of Target Species Captured at the Breitenbush River To-Date

\*Fish that were missing heads are not included in length and weight calculations.

| May 1-15, 2023       |       |         |       |           |              |     |      |     |                         |      |  |
|----------------------|-------|---------|-------|-----------|--------------|-----|------|-----|-------------------------|------|--|
| Site                 | Route | Species | Life  | Collected | Length (mm)* |     |      |     | Weight (g) <sup>*</sup> |      |  |
|                      |       |         | stage |           | Min          | Max | Mean | Min | Max                     | Mean |  |
|                      |       | CHS     | Fry   | 0         | N/A          | N/A | N/A  | N/A | N/A                     | N/A  |  |
|                      |       | CHS     | Parr  | 0         | N/A          | N/A | N/A  | N/A | N/A                     | N/A  |  |
| Breitenbush<br>River | 5ft   | CHS     | Smolt | 0         | N/A          | N/A | N/A  | N/A | N/A                     | N/A  |  |
|                      |       | STW     | Fry   | 0         | N/A          | N/A | N/A  | N/A | N/A                     | N/A  |  |
|                      |       | STW     | Parr  | 0         | N/A          | N/A | N/A  | N/A | N/A                     | N/A  |  |
|                      |       | STW     | Smolt | 0         | N/A          | N/A | N/A  | N/A | N/A                     | N/A  |  |

\*Fish that were missing heads are not included in length and weight calculations.

#### **Trapping Efficiency**

No trapping efficiency trials have been performed at this site to date.

#### Injuries and Copepod Infection

No target species have been captured at this site to date. A summary of injury data for fish captured during this reporting period are shown in table 5.

# Table 5. Number of Descaled, Bodily/Eye Injured, Copepod Infected and Dead Chinook Salmon and Winter Steelhead for Sampling Period (Breitenbush River).

| Site                 | Species             | # Fish<br>Collected | # DSC*<br><20% | # DSC*<br>>20% | # with<br>Body<br>Injuries | # with<br>Eye<br>Injuries | # with<br>COP*<br>In B.C. | # with<br>COP*<br>on Fins | Mortalities |
|----------------------|---------------------|---------------------|----------------|----------------|----------------------------|---------------------------|---------------------------|---------------------------|-------------|
| Breitenbush<br>River | Chinook             | 0                   | 0              | 0              | 0                          | 0                         | 0                         | 0                         | 0           |
|                      | Winter<br>Steelhead | 0                   | 0              | 0              | 0                          | 0                         | 0                         | 0                         | 0           |

\*DSC=Descaled, COP=Copepods, B.C.=Branchial Cavity

#### Collected DNA and Scale Samples

No target fish were captured during this reporting period.

#### **PIT Tags**

No fish were PIT tagged during this reporting period.

#### **Non-Target Species**

No non-target species were captured during this reporting period. A summary of non-target fish capture is provided in table 6.

| Species         | 5 ft<br>Capture | 5 ft<br>Mortality | Season<br>Total | Season Total<br>Mortality |  |  |  |  |  |  |
|-----------------|-----------------|-------------------|-----------------|---------------------------|--|--|--|--|--|--|
| Kokanee         | 0               | 0                 | 0               | 0                         |  |  |  |  |  |  |
| Cutthroat Trout | 0               | 0                 | 0               | 0                         |  |  |  |  |  |  |
| Sculpin         | 0               | 0                 | 0               | 0                         |  |  |  |  |  |  |
| Totals          | 0               | 0                 | 0               | 0                         |  |  |  |  |  |  |

#### Table 6. Summary of Non-target Species (Breitenbush River).

#### **Stream Statistics**

Basic stream statistics at the proposed Breitenbush River RST site were calculated from data downloaded from the U.S. Geological Survey stream gauge number 14179000. Instantaneous discharge (cfs) and Gauge height (feet) flow metrics are available at this gauge. During the reporting period, daily maximum values for instantaneous discharge ranged from 1,050.0 cfs to 1,930.0 (mean: 1,526.0 cfs). Figure 7 shows instantaneous discharge.

Stream temperatures will be recorded every 2 hours for the length of the reporting period for the RST (Figure 8).

Catch per unit of effort (CPUE) data are summarized in Table 7. Gage height and capture data for the duration of monitoring efforts at this location are provided in Appendix B.

|                | Chinook | Winter Steelhead |
|----------------|---------|------------------|
| Description    | (5 ft)  | (5 ft)           |
| Catch          | 0       | 0                |
| Effort (hrs)   | 0       | 0                |
| CPUE (fish/hr) | 0       | 0                |

Table 7. Summary of salmonid CPUE, Breitenbush River.





Figure 7. Discharge (cfs); Breitenbush River



#### Figure 8. Temperature at RST (Breitenbush River)

\*Note: Temperature data was supplemented from USGS gage 14179000 as the trap is not yet deployed nor is a temperature probe.

#### North Santiam River – Detroit Head of Reservoir

The Detroit Head of Reservoir- North Santiam River RST was installed on April 19<sup>th</sup>, 2023. This site started sampling on May 4, 2023.

#### **Target Species**

This reporting period began on May 1<sup>st</sup> and ended on May 15<sup>th</sup>. There were a total of 2,098 Chinook Salmon (CHS) and 114 Winter Steelhead (STW) captured during the 15-day sampling period (Figure 9). Sampling duration was 80.0% of the reporting period for the RST. Figure 10 shows length frequency data to-date. Table 8 provides life stage, length, and weight data for all Chinook Salmon and Winter Steelhead that have been caught at the Detroit Head of Reservoir site to-date and for the reporting period.




Figure 9. Chinook and Winter Steelhead Captured per day 05/01/2023 to 05/15/2023 (Detroit Head of Reservoir)




Figure 10. Length Frequency of Juvenile Chinook and Winter Steelhead Sampled Season To-Date (Detroit Head of Reservoir)

| Table 8. Descriptive Statistics of Target Species Captured at Detroit Head of ReservoirSeason To-Date | , |
|---|---|
| To-Date (Since May 04, 2023)  |   |

|                | I O-Date (Since May 04, 2023)             |     |       |       |     |     |      |                         |     |      |  |  |
|----------------|---|-----|-------|-------|-----|-----|------|-------------------------|-----|------|--|--|
| Site           | Route Species Life Collected Length (mm)* |     |       |       |     |     |      | Weight (g) <sup>*</sup> |     |      |  |  |
|                |   |     | stage |       | Min | Мах | Mean | Min                     | Мах | Mean |  |  |
|                |   | CHS | Fry   | 2,098 | 29  | 45  | 34.7 | N/A                     | N/A | N/A  |  |  |
|                |   | CHS | Parr  | 0     | N/A | N/A | N/A  | N/A                     | N/A | N/A  |  |  |
| Detroit<br>HOR | 5ft                                       | CHS | Smolt | 0     | N/A | N/A | N/A  | N/A                     | N/A | N/A  |  |  |
|                |   | STW | Fry   | 114   | 30  | 39  | 34.6 | N/A                     | N/A | N/A  |  |  |
|                |   | STW | Parr  | 0     | N/A | N/A | N/A  | N/A                     | N/A | N/A  |  |  |
|                |   | STW | Smolt | 0     | N/A | N/A | N/A  | N/A                     | N/A | N/A  |  |  |

\*Fish that were missing heads are not included in length and weight calculations.

|            | May 1-15, 2023 |         |                       |         |         |         |      |           |     |          |      |  |           |    |
|------------|----------------|---------|-----------------------|---------|---------|---------|------|-----------|-----|----------|------|--|-----------|----|
| Site Route |                | Species | Species Life<br>stage | Species | Species | Species |      | Collected | L   | ength (m | ım)* |  | Weight (g | )* |
|            |                | stage   |                       |         | Min     | Мах     | Mean | Min       | Мах | Mean     |      |  |           |    |
|            |                | CHS     | Fry                   | 2,098   | 29      | 45      | 34.7 | N/A       | N/A | N/A      |      |  |           |    |
|            |                | CHS     | Parr                  | 0       | N/A     | N/A     | N/A  | N/A       | N/A | N/A      |      |  |           |    |
| Detroit    | 5ft            | CHS     | Smolt                 | 0       | N/A     | N/A     | N/A  | N/A       | N/A | N/A      |      |  |           |    |
| HOR        |                | STW     | Fry                   | 114     | 30      | 39      | 34.6 | N/A       | N/A | N/A      |      |  |           |    |
|            |                | STW     | Parr                  | 0       | N/A     | N/A     | N/A  | N/A       | N/A | N/A      |      |  |           |    |
|            |                | STW     | Smolt                 | 0       | N/A     | N/A     | N/A  | N/A       | N/A | N/A      |      |  |           |    |

\*Fish that were missing heads are not included in length and weight calculations.

# **Trapping Efficiency**

Trapping Efficiency has yet to be conducted at the Detroit Head of Reservoir RST since its installation.

#### Injuries and Copepod Infection

Partial descaling <20% was observed in 1 of the 2,098 Chinook captured (0.04%), 0 displayed descaling >20% (0.0%), 33 displayed body injury (1.5%), 4 had eye injury (0.19%), 0 had copepods present in the branchial cavity (0.0%) and 0 had copepods on fins (0.0%). 0 Chinook displayed gas bubble disease (0.0%). There were 5 mortalities (0.24%).

Partial descaling <20% was observed on 0 of the 114 Winter Steelhead captured (0.0%) and 0 displayed descaling >20% (0.0%), 2 displayed body injury (1.75%), 1 had eye injury (0.87%), 0 had copepods present in the branchial cavity (0.0%) and 0 had copepods on fins (0.0%). 0 Winter Steelhead displayed gas bubble disease (0.0%). There were 0 mortalities (0.0%). Injury data is summarized in 8.

#### Table 9. Number of Descaled, Bodily/Eye Injured, Copepod Infected and Dead Chinook Salmon and Winter Steelhead for Sampling Period (Detroit Head of Reservoir).

| Site           | Species             | # Fish<br>Collected | #<br>DSC*<br><20% | #<br>DSC*<br>>20% | # with<br>Body<br>Injuries | # with<br>Eye<br>Injuries | # with<br>COP*<br>In B.C. | # with<br>COP*<br>on Fins | Mortalities |
|----------------|---------------------|---------------------|-------------------|-------------------|----------------------------|---------------------------|---------------------------|---------------------------|-------------|
| Detroit<br>HOR | Chinook             | 2,098               | 1                 | 1                 | 33                         | 4                         | 0                         | 0                         | 5           |
| HUK .          | Winter<br>Steelhead | 114                 | 0                 | 0                 | 2                          | 1                         | 0                         | 0                         | 0           |

\*DSC=Descaled, COP=Copepods, B.C.=Branchial Cavity

#### Collected DNA and Scale Samples

For the reporting period, DNA was collected from 1 Spring Chinook and 0 Winter Steelhead. The other targets captured did not meet length criteria for scales or DNA.

#### **PIT Tags**

No fish caught in the RST have met the length criteria for PIT tagging this reporting period.

#### **VIE Marking**

Visual Implant Elastomer (VIE) trials commenced at the Detroit Head of Reservoir – North Santiam River site on 5/5/2023. VIE tag color and locations are changed every month to distinctly mark groups of fish by capture date. Since then, 889 Chinook and 60 Winter Steelhead have been VIE marked with fluorescent elastomer. No fish with VIE marks have been detected at downstream RST sites to date. Fish still showing an egg sac are not VIE marked.

| Date Tagged         | Species   | Tag Location | VIE Color | # Tagged | # Recaptured to<br>Date |
|---------------------|-----------|--------------|-----------|----------|-------------------------|
| 5/01/2023-5/15/2023 | Chinook   | Right Dorsal | Orange    | 889      | 0                       |
| 5/01/2023-5/15/2023 | O. mykiss | Right Dorsal | Orange    | 60       | 0                       |

#### **Non-Target Species**

15 non-target species fish were captured during the reporting period; the data is summarized below in 10.

| Species         | 5 ft<br>Capture | 5 ft<br>Mortality | Season<br>Total | Season Total<br>Mortality |
|-----------------|-----------------|-------------------|-----------------|---------------------------|
| Kokanee         | 13              | 0                 | 13              | 0                         |
| Cutthroat Trout | 1               | 0                 | 1               | 0                         |
| Sculpin         | 1               | 0                 | 1               | 0                         |
| Totals          | 15              | 0                 | 15              | 0                         |

#### Table 10. Summary of Non-target Species (Detroit Head of Reservoir).

#### **Stream Statistics**

Basic stream statistics at the Detroit Head of Reservoir site were calculated from data downloaded from U.S. Geological Survey stream gauge number 14178000. Gauge height (feet) and Discharge (cfs) metrics are provided at gauge 14178000. During the reporting period, daily maximum values for instantaneous discharge ranged from 1,920.0 cfs to 3,090.0 cfs (mean: 2,547.3 cfs) during the reporting period. Figure 11 shows instantaneous discharge.

Stream temperatures were recorded every 2 hours for the length of the reporting period at the Detroit Head of Reservoir RST site. Figure 12 shows temperature during the reporting period.

Catch per unit of effort (CPUE) data are summarized in Table 11. Discharge and capture data for the duration of monitoring efforts at this location are provided in Appendix B.

|                | Chinook | Winter Steelhead |
|----------------|---------|------------------|
| Description    | (5 ft)  | (5 ft)           |
| Catch          | 2,098   | 114              |
| Effort (hrs)   | 265.0   | 265.0            |
| CPUE (fish/hr) | 7.917   | 0.430            |

Table 11. Summary of salmonid CPUE, Detroit Head of Reservoir – North Santiam River.




Figure 11. Discharge (cfs); Detroit Head of Reservoir – North Santiam River




Figure 12. Temperature at RST (Detroit Head of Reservoir – North Santiam River)

## Middle Santiam River- Green Peter Head of Reservoir

The Green Peter Head of Reservoir- Middle Santiam River RST was installed on April 26<sup>th</sup>, 2023. This site started sampling on May 4<sup>th</sup>, 2023. All natural origin O. mykiss captured at this site will be reported as Winter Steelhead.

#### **Target Species**

This reporting period began on May 1<sup>st</sup> and ended on May 15<sup>th</sup>. There were a total of 20 Chinook Salmon (CHS) and 1 Winter Steelhead (STW) captured during the 15-day sampling period (Figure 13). Sampling duration was 80.0% of the reporting period for the RST. Figure 14 shows length frequency data to-date. Table 12 provides life stage, length, and weight data for all Chinook Salmon and Winter Steelhead that have been caught at the Middle Santiam River- Green Peter Head of Reservoir site to-date and for the reporting period.




Figure 13. Chinook Captured per day 5/01/2023 to 5/15/2023 (Green Peter Head of Reservoir – Middle Santiam River)




Figure 14. Length Frequency of Juvenile Chinook Sampled Season To-Date (Green Peter Head of Reservoir – Middle Santiam River)

|                    | To-date (since May 4, 2023) |         |       |           |     |           |      |     |                         |      |  |  |  |
|--------------------|-----------------------------|---------|-------|-----------|-----|-----------|------|-----|-------------------------|------|--|--|--|
| Site               | Route                       | Species | Life  | Collected | L   | .ength (m | m)*  |     | Weight (g) <sup>*</sup> |      |  |  |  |
|                    |                             |         | stage |           | Min | Мах       | Mean | Min | Мах                     | Mean |  |  |  |
|                    |                             | CHS     | Fry   | 20        | 33  | 45        | 36.3 | N/A | N/A                     | N/A  |  |  |  |
| Green              | 5ft                         | CHS     | Parr  | 0         | N/A | N/A       | N/A  | N/A | N/A                     | N/A  |  |  |  |
| Peter<br>Head of   |                             | CHS     | Smolt | 0         | N/A | N/A       | N/A  | N/A | N/A                     | N/A  |  |  |  |
| Reservoir          |                             | STW     | Fry   | 1         | 36  | 36        | 36   | N/A | N/A                     | N/A  |  |  |  |
| -Middle<br>Santiam |                             | STW     | Parr  | 0         | N/A | N/A       | N/A  | N/A | N/A                     | N/A  |  |  |  |
|                    |                             | STW     | Smolt | 0         | N/A | N/A       | N/A  | N/A | N/A                     | N/A  |  |  |  |

#### Table 12. Descriptive Statistics of Target Species Captured at Green Peter Head of Reservoir – Middle Santiam River Season To-Date

\*Fish that were missing heads are not included in length and weight calculations.

|                      | May 1-15, 2023 |              |               |           |     |           |      |     |                         |      |  |  |  |
|----------------------|----------------|--------------|---------------|-----------|-----|-----------|------|-----|-------------------------|------|--|--|--|
| Site                 | Route          | oute Species | Life<br>stage | Collected | L   | .ength (m | m)*  |     | Weight (g) <sup>*</sup> |      |  |  |  |
|                      |                |              |               |           | Min | Мах       | Mean | Min | Мах                     | Mean |  |  |  |
|                      |                | CHS          | Fry           | 20        | 33  | 45        | 36.3 | N/A | N/A                     | N/A  |  |  |  |
| Green                |                | CHS          | Parr          | 0         | N/A | N/A       | N/A  | N/A | N/A                     | N/A  |  |  |  |
| Peter                | 5ft            | CHS          | Smolt         | 0         | N/A | N/A       | N/A  | N/A | N/A                     | N/A  |  |  |  |
| Head of<br>Reservoir |                | STW          | Fry           | 1         | 36  | 36        | 36   | N/A | N/A                     | N/A  |  |  |  |
| -Middle<br>Santiam   |                | STW          | Parr          | 0         | N/A | N/A       | N/A  | N/A | N/A                     | N/A  |  |  |  |
|                      |                | STW          | Smolt         | 0         | N/A | N/A       | N/A  | N/A | N/A                     | N/A  |  |  |  |

\*Fish that were missing heads are not included in length and weight calculations.

## **Trapping Efficiency**

Trapping Efficiency has yet to be conducted at the Green Peter Head of Reservoir – Middle Santiam River RST since it began fishing on May 4<sup>th</sup>, 2023.

#### Injuries and Copepod Infection

Partial descaling <20% was observed in 0 of the 20 Chinook captured (0.0%), 0 displayed descaling >20% (0.0%), 1 displayed body injury (5.0%), 0 had eye injuries (0.0%), 0 had copepods present in the branchial cavity (0.0%) and 0 had copepods on fins (0.0%). 0 Chinook displayed gas bubble disease (0.0%). There were 0 mortalities (0.0%).

Partial descaling <20% was observed on 0 of the 1 Winter Steelhead captured (0.0%) and 0 displayed descaling >20% (0.0%), 0 displayed body injury (0.0%), 0 had eye injuries (0.0%), 0 had copepods present in the branchial cavity (0.0%) and 0 had copepods on fins (0.0%). 0 Winter Steelhead displayed gas bubble disease (0.0%). There were 0 mortalities (0.0%). Injury data is summarized in Table 13.

# Table 13. Number of Descaled, Bodily/Eye Injured, Copepod Infected and Dead ChinookSalmon and Winter Steelhead for Sampling Period (Green Peter Head of Reservoir-MiddleSantiam River).

| Site                            | Species             | # Fish<br>Collected | # DSC*<br><20% | # DSC*<br>>20% | # with<br>Body<br>Injuries | # with<br>Eye<br>Injuries | # with<br>COP*<br>In B.C. | # with<br>COP*<br>on<br>Fins | Mortalities |
|---------------------------------|---------------------|---------------------|----------------|----------------|----------------------------|---------------------------|---------------------------|------------------------------|-------------|
| Green<br>Peter Head<br>of       | Chinook             | 20                  | 0              | 0              | 1                          | 0                         | 0                         | 0                            | 0           |
| Reservoir-<br>Middle<br>Santiam | Winter<br>Steelhead | 1                   | 0              | 0              | 0                          | 0                         | 0                         | 0                            | 0           |

\*DSC=Descaled, COP=Copepods, B.C.=Branchial Cavity

#### Collected DNA and Scale Samples

For the reporting period, DNA was collected from 1 Spring Chinook and 0 Winter Steelhead. The other targets captured did not meet length criteria for DNA sampling or were too descaled/damaged to collect samples.

#### **PIT Tags**

No Spring Chinook or Winter Steelhead were PIT tagged during this reporting period. All fish captured did not meet the size criteria for PIT tagging. More information regarding PIT tagged fish can be found in Appendix D.

#### VIE Marking

Visual Implant Elastomer (VIE) trials commenced at the Green Peter Head of Reservoir – Middle Santiam River site on 5/5/2023. VIE tag color and locations are changed every month to distinctly mark groups of fish by capture date. Since then, 14 Chinook and 1 Winter Steelhead have been VIE marked with

fluorescent elastomer. No fish with VIE marks have been detected at downstream RST sites to date. Fish still showing an egg sac are not VIE marked.

| Date Tagged         | Species   | Tag<br>Location | VIE Color | # Tagged | # Recaptured to<br>Date |
|---------------------|-----------|-----------------|-----------|----------|-------------------------|
| 5/01/2023-5/15/2023 | Chinook   | Right Dorsal    | Orange    | 14       | 0                       |
| 5/01/2023-5/15/2023 | O. mykiss | Right Dorsal    | Orange    | 1        | 0                       |

#### **Non-Target Species**

5 non-target fish were collected during the reporting period; the data is summarized below in Table 14.

# Table 14. Summary of Non-target Species (Green Peter Head of Reservoir – Middle Santiam River).

| Species         | 5 ft 5 ft<br>Capture Mortality |   | Season<br>Total | Season Total<br>Mortality |
|-----------------|--------------------------------|---|-----------------|---------------------------|
| Kokanee         | 5                              | 0 | 5               | 0                         |
| Cutthroat Trout | 0                              | 0 | 0               | 0                         |
| Sculpin         | 0                              | 0 | 0               | 0                         |
| Totals          | 5                              | 0 | 5               | 0                         |

#### **Stream Statistics**

Basic stream statistics at the Green Peter Head of Reservoir – Middle Santiam River site were calculated from data downloaded from the U.S. Geological Survey stream gauge number 14185800. Gauge height (feet) is the only flow metric available at this gauge. During the reporting period, daily maximum values for gage height ranged from 4.0 ft to 5.2 ft (mean: 4.5 ft). Figure 15 shows gage height.

Stream temperatures were recorded every 2 hours for the length of the report period for the RST (Figure 16). Temperature probes for the trap operated normally throughout this reporting period.

Catch per unit of effort (CPUE) data are summarized in Table 15. Gage height and capture data for the duration of monitoring efforts at this location are provided in Appendix B.

|                | Chinook | Winter Steelhead |
|----------------|---------|------------------|
| Description    | (5 ft)  | (5 ft)           |
| Catch          | 20      | 1                |
| Effort (hrs)   | 261.4   | 261.4            |
| CPUE (fish/hr) | 0.077   | 0.004            |

 Table 15. Summary of salmonid CPUE, Green Peter Head of Reservoir – Middle Santiam River.



Figure 15. Gage Height (feet); Green Peter Head of Reservoir – Middle Santiam River




Figure 16. Temperature at RST (Green Peter Head of Reservoir – Middle Santiam River)

## Middle Fork Willamette River- Hills Creek Head of Reservoir

The Hills Creek Head of Reservoir RST was installed and began sampling on May 9th, 2023.

#### **Target Species**

This reporting period began on May 1<sup>st</sup> and ended on May 15<sup>th</sup>. There were a total of 25 Chinook Salmon captured during the 15-day sampling period (Figure 17). Sampling duration was 46.7% of the reporting period for the RST. Figure 18 shows length frequency data to-date. Table 16 provides life stage, length, and weight data for all Chinook Salmon that have been caught at the Middle Fork Willamette River- Hills Creek Head of Reservoir site to-date and for the reporting period.




Figure 17. Chinook Captured per day 05/01/2023 to 05/15/2023 (Hills Creek Head of Reservoir)




Figure 18. Length Frequency of Juvenile Chinook Sampled Season To-Date (Hills Creek Head of Reservoir)

#### Table 16. Descriptive Statistics of Target Species Captured at Hills Creek Head of Reservoir Season To-Date

|                      | To-Date (Since May 9 <sup>th</sup> , 2023) |         |               |           |                          |     |      |                         |     |      |  |  |  |
|----------------------|--|---------|---------------|-----------|--------------------------|-----|------|-------------------------|-----|------|--|--|--|
| Site                 | Route                                      | Species | Life<br>stage | Collected | Length (mm) <sup>.</sup> |     |      | Weight (g) <sup>.</sup> |     |      |  |  |  |
|                      |  |         |               |           | Min                      | Мах | Mean | Min                     | Мах | Mean |  |  |  |
| Hills Creek          |  | CHS     | Fry           | 24        | 32                       | 50  | 36.6 | N/A                     | N/A | N/A  |  |  |  |
| Head of<br>Reservoir | 5 ft                                       | CHS     | Parr          | 1         | 57                       | 57  | 57   | 1.9                     | 1.9 | 1.9  |  |  |  |
|                      |  | CHS     | Smolt         | 0         | N/A                      | N/A | N/A  | N/A                     | N/A | N/A  |  |  |  |

\*Most fry are too small to collect accurate weights and thus some metrics are not available for them.

|                      | May 1-15, 2023 |         |               |           |  |     |      |     |     |                  |  |  |  |
|----------------------|----------------|---------|---------------|-----------|--|-----|------|-----|-----|------------------|--|--|--|
| Site                 | Route          | Species | Life<br>stage | Collected | Length (mm) <sup>.</sup> Weight (g) <sup>.</sup> |     |      |     |     | (g) <sup>.</sup> |  |  |  |
|                      |                |         |               |           | Min  | Мах | Mean | Min | Max | Mean             |  |  |  |
| Hills Creek          |                | CHS     | Fry           | 24        | 32   | 50  | 36.6 | N/A | N/A | N/A              |  |  |  |
| Head of<br>Reservoir | 5 ft           | CHS     | Parr          | 1         | 57   | 57  | 57   | 1.9 | 1.9 | 1.9              |  |  |  |
|                      |                | CHS     | Smolt         | 0         | N/A  | N/A | N/A  | N/A | N/A | N/A              |  |  |  |

\*Most fry are too small to collect accurate weights and thus some metrics are not available for them.

#### **Trapping Efficiency**

Trapping Efficiency was not conducted during the reporting period since it began fishing on May 9th, 2023.

#### Injuries and Copepod Infection

25 Chinook were captured for the reporting period. Of the fish captured, partial descaling <20% was observed on 1 fish (4.0%) and descaling >20% was observed on 0 fish (0.0%). 0 fish had bodily injury (0.0%). 0 fish displayed eye injuries (0.0%). 0 fish had copepods in the branchial cavity (0.0%), 0 had copepods on fins (0.0%). There were 0 mortalities for this reporting period (0.0%). Injury data for the reporting period is summarized in Table 17. To date injury data can be found in Appendix A.

#### Table 17. Number of Descaled, Bodily/Eye Injured, Copepod Infected and Dead Chinook Salmon and Winter Steelhead for Sampling Period (Hills Creek Head of Reservoir).

| Site                                | # CHS<br>Collected | # DSC*<br><20% | # DSC*<br>>20% | # with<br>Body<br>Injuries | # with Eye<br>Injuries | # with COP* | # with COP* | Mortalities |
|-------------------------------------|--------------------|----------------|----------------|----------------------------|------------------------|-------------|-------------|-------------|
| Hills Creek<br>Head of<br>Reservoir | 25                 | 1              | 0              | 0                          | 0                      | 0           | 0           | 0           |

\*DSC=Descaled, COP=Copepods, B.C.=Branchial Cavity

#### Collected DNA and Scale Samples

DNA was collected from 4 of the Chinook captured and scales were collected from 2 of the Chinook captured. The rest of the captured fish were under the minimum fork length threshold and samples were not collected.

#### **PIT Tags**

No Spring Chinook were PIT tagged during this reporting period. The fish captured did not meet the size criteria for PIT tagging. More information regarding PIT tagged fish can be found in Appendix D.

#### VIE Marking

Visual Implant Elastomer (VIE) trials commenced at the Hills Creek Head of Reservoir site on 5/10/2023. VIE tag color and locations are changed every month to distinctly mark groups of fish by capture date. Since then, 15 Chinook have been VIE marked with fluorescent elastomer. No fish with VIE marks have been detected at downstream RST sites to date. Fish still showing an egg sac are not VIE marked.

| Date Tagged        | Tag Location | VIE Color | # Tagged | # Recaptured to Date |
|--------------------|--------------|-----------|----------|----------------------|
| 5/1/2023-5/15/2023 | Left Dorsal* | Orange    | 6        | 0                    |
| 5/1/2023-5/15/2023 | Right Dorsal | Orange    | 9        | 0                    |

\*denotes fish marked at incorrect location for that period.

#### **Non-Target Species**

55 non-target species fish were captured during the reporting period; the data is summarized below in Table 18.

|                   | <u> </u>        | -                 |                 |                           |
|-------------------|-----------------|-------------------|-----------------|---------------------------|
| Species           | 5 ft<br>Capture | 5 ft<br>Mortality | Season<br>Total | Season Total<br>Mortality |
| Dace              | 16              | 0                 | 16              | 0                         |
| Cutthroat Trout   | 1               | 0                 | 1               | 0                         |
| Sculpin           | 4               | 1                 | 4               | 1                         |
| Largescale Sucker | 28              | 1                 | 28              | 1                         |
| Redside Shiner    | 6               | 0                 | 6               | 0                         |
| Totals            | 55              | 2                 | 55              | 2                         |

#### Table 18. Summary of Non-target Species (Hills Creek Head of Reservoir).

#### **Stream Statistics**

Basic stream statistics at the Hills Creek Head of Reservoir site were calculated from data downloaded from the U.S. Geological Survey stream gauge number 14144800. Gauge height (feet) is the only flow metric available at this gauge. During the reporting period, daily maximum values for gage height ranged from 10.2 ft to 12.0 ft (mean: 10.9 ft). Figure 20 shows gage height.

Stream temperatures were recorded every 2 hours for the length of the report period for the RST (Figure 21). Temperature probes for the trap operated normally throughout this reporting period.

Catch per unit of effort (CPUE) data are summarized in Table 19. Gage height and capture data for the duration of monitoring efforts at this location are provided in Appendix B.

|                | Chinook |
|----------------|---------|
| Description    | 5 ft    |
| Catch          | 25      |
| Effort (hrs)   | 139.6   |
| CPUE (fish/hr) | 0.179   |

#### Table 19. Summary of salmonid CPUE, Hills Creek Head of Reservoir.





Figure 20. Gage Height (feet); Hills Creek Head of Reservoir, Near Oakridge, OR



#### Figure 21. Temperature at RST (Hills Creek Head of Reservoir)

\*Note: Trap was installed 5/9/2023 and the temperature logger started recording on that date, temperature for the reporting period is supplemented with USGS stream gage 14144800

#### **Issues Encountered**

None.

# Upcoming USACE Support Services

None at this time.

# Appendix A Chinook (CHS) To-Date

|                 |               |      |      |     |     | Cł  | ninod | ok In | jurie | s to- | date |     |    |    |     |     |     |     |     |     |     |     |     |
|-----------------|---------------|------|------|-----|-----|-----|-------|-------|-------|-------|------|-----|----|----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| p/Life Stage    | Total<br>Fish | MUNK | DS<2 | BLO | EYB | FUN | BKD   | сор   | DS>2  | PRD   | FID  | НВО | BO | ЮН | BVT | НВР | BRU | TEA | ОРD | HIN | FVB | РОР | GBD |
| Detroit HOR     | 2098          |      |      |     |     |     |       |       |       |       |      |     |    |    |     |     |     |     |     |     |     |     |     |
| 5 ft            | 2098          | 1    | 1    |     | 4   |     |       |       | 1     |       | 4    |     |    |    |     |     | 8   | 5   | 15  | 5   | 17  |     |     |
| Fry             | 2098          | 1    | 1    |     | 4   |     |       |       | 1     |       | 4    |     |    |    |     |     | 8   | 5   | 15  | 5   | 17  |     |     |
| Green Peter HOR |               |      |      |     |     |     |       |       |       |       |      |     |    |    |     |     |     |     |     |     |     |     |     |
| 5 ft            | 20            |      |      |     |     |     |       |       |       |       | 1    |     |    |    |     |     | 1   |     |     |     |     |     |     |
| Fry             | 20            |      |      |     |     |     |       |       |       |       | 1    |     |    |    |     |     | 1   |     |     |     |     |     |     |
| Hills Creek HOR |               |      |      |     |     |     |       |       |       |       |      |     |    |    |     |     |     |     |     |     |     |     |     |
| 5 ft            | 25            |      | 1    |     |     |     |       |       |       |       |      |     |    |    |     |     |     |     |     |     |     |     |     |
| Fry             | 24            |      | 1    |     |     |     |       |       |       |       |      |     |    |    |     |     |     |     |     |     |     |     |     |
| Parr            | 1             |      |      |     |     |     |       |       |       |       |      |     |    |    |     |     |     |     |     |     |     |     |     |

# Chinook (CHS) During Reporting Period

|                 | (             | Chino | ok I | njuri | es D | urin | g Re | porti | ing P | erio | d (5- | 01-2 | 023 t | : <b>o 5</b> -1 | 15-20 | )23) |     |     |     |     |     |     |     |
|-----------------|---------------|-------|------|-------|------|------|------|-------|-------|------|-------|------|-------|-----------------|-------|------|-----|-----|-----|-----|-----|-----|-----|
| p/Life Stage    | Total<br>Fish | MUNK  | DS<2 | BLO   | ЕYВ  | FUN  | BKD  | сор   | DS>2  | PRD  | FID   | НВО  | BO    | ОН              | BVT   | НВР  | BRU | TEA | ОРD | NIN | FVB | РОР | GBD |
| Detroit HOR     | 2098          |       |      |       |      |      |      |       |       |      |       |      |       |                 |       |      |     |     |     |     |     |     |     |
| 5 ft            | 2098          | 1     | 1    |       | 4    |      |      |       | 1     |      | 4     |      |       |                 |       |      | 8   | 5   | 15  | 5   | 17  |     |     |
| Fry             | 2098          | 1     | 1    |       | 4    |      |      |       | 1     |      | 4     |      |       |                 |       |      | 8   | 5   | 15  | 5   | 17  |     |     |
| Green Peter HOR |               |       |      |       |      |      |      |       |       |      |       |      |       |                 |       |      |     |     |     |     |     |     |     |
| 5 ft            | 20            |       |      |       |      |      |      |       |       |      | 1     |      |       |                 |       |      | 1   |     |     |     |     |     |     |
| Fry             | 20            |       |      |       |      |      |      |       |       |      | 1     |      |       |                 |       |      | 1   |     |     |     |     |     |     |
| Hills Creek HOR |               |       |      |       |      |      |      |       |       |      |       |      |       |                 |       |      |     |     |     |     |     |     |     |
| 5 ft            | 25            |       | 1    |       |      |      |      |       |       |      |       |      |       |                 |       |      |     |     |     |     |     |     |     |
| Fry             | 24            |       | 1    |       |      |      |      |       |       |      |       |      |       |                 |       |      |     |     |     |     |     |     |     |
| Parr            | 1             |       |      |       |      |      |      |       |       |      |       |      |       |                 |       |      |     |     |     |     |     |     |     |

# Steelhead (O. mykiss) To Date


|                 |               |      |             |     |     | тук | iss ir | njurie | es to | -date | e   |    |   |     | _   |     |     |     |             |     |     |     |
|-----------------|---------------|------|-------------|-----|-----|-----|--------|--------|-------|-------|-----|----|---|-----|-----|-----|-----|-----|-------------|-----|-----|-----|
| p/Life Stage    | Total<br>Fish | MUNK | DS<2<br>BLO | EYB | FUN | BKD | СОР    | DS>2   | PRD   | FID   | НВО | BO | 오 | BVT | НВР | BRU | TEA | OPD | N<br>N<br>H | FVB | POP | GBD |
| Detroit HOR     |               |      |             |     |     |     |        |        |       |       |     |    |   |     |     |     |     |     |             |     |     |     |
| 5 ft            | 114           |      |             | 1   |     |     |        |        |       |       |     |    |   | 1   |     |     |     | 1   |             | 1   |     |     |
| Fry             | 114           |      |             | 1   |     |     |        |        |       |       |     |    |   | 1   |     |     |     | 1   |             | 1   |     |     |
| Green Peter HOR |               |      |             |     |     |     |        |        |       |       |     |    |   |     |     |     |     |     |             |     |     |     |
| 5 ft            | 1             |      |             |     |     |     |        |        |       |       |     |    |   |     |     |     |     |     |             |     |     |     |
| Fry             | 1             |      |             |     |     |     |        |        |       |       |     |    |   |     |     |     |     |     |             |     |     |     |

|                 | O. mykiss Injuries During Reporting Period (5-01-2023 to 5-15-2023) |                           |            |             |            |          |                    |                   |                   |  |
|-----------------|---|---------------------------|------------|-------------|------------|----------|--------------------|-------------------|-------------------|--|
| p/Life Stage    | Total<br>Fish   | DS<2<br>BLO<br>EYB<br>FUN | BKD<br>COP | DS>2<br>PRD | FID<br>HBO | BO<br>HO | BV I<br>HBP<br>BRU | TEA<br>OPD<br>HIN | FVB<br>POP<br>GBD |  |
| Detroit HOR     |   |                           |            |             |            |          |                    |                   | 1                 |  |
| 5 ft            | 114   | 1                         |            |             |            |          | 1                  | 1                 | 1                 |  |
| Fry             | 114   | 1                         |            |             |            |          | 1                  | 1                 | 1                 |  |
| Green Peter HOR |   |                           |            |             |            |          |                    |                   |                   |  |
| 5 ft            | 1   |                           |            |             |            |          |                    |                   |                   |  |
| Fry             | 1   |                           |            |             |            |          |                    |                   |                   |  |

# Steelhead (O. mykiss) During Reporting Period

Appendix B Detroit Head of Reservoir-North Santiam River Operational and Capture Data Since Start of Monitoring







#### Green Peter Head of Reservoir-Middle Santiam River Operational and Capture Data Since Start of Monitoring





#### Hills Creek Head of Reservoir-Middle Fork Willamette River Operational and Capture Data Since Start of Monitoring





#### Appendix C

#### Summary of PIT Tagged Fish for Reporting Period

| Site | Тгар | # of PIT Tagged Fish |
|------|------|----------------------|
| N/A  | N/A  | N/A                  |

#### Summary of Captured Fish Containing PIT Tags This Season

| Site | Тгар | # of PIT Tagged Fish |
|------|------|----------------------|
| N/A  | N/A  | N/A                  |

#### Summary of EAS VIE Marked Fish for Reporting Period

| •  |      |               |           |       |
|--|------|---------------|-----------|-------|
| Site   | Trap | VIE Mark Code | Species   | # VIE |
| Detroit Head of Reservoir – North Santiam River      | 5 ft | RDO           | Chinook   | 889   |
| Detroit Head of Reservoir – North Santiam River      | 5 ft | RDO           | O. mykiss | 60    |
| Green Peter Head of Reservoir – Middle Santiam River | 5 ft | RDO           | Chinook   | 14    |
| Green Peter Head of Reservoir – Middle Santiam River | 5 ft | RDO           | O. mykiss | 1     |
| Hills Creek Head of Reservoir                        | 5 ft | RDO           | Chinook   | 9     |
| Hills Creek Head of Reservoir                        | 5 ft | LDO*          | Chinook   | 6     |

RDO denotes location and color (Right Dorsal Orange)

\*6 fish were accidentally tagged in the wrong location

#### List of EAS PIT Tagged Fish for Reporting Period

| Site | Тгар | PIT Tag | Date | Species |
|------|------|---------|------|---------|
| N/A  | N/A  | N/A     | N/A  | N/A     |