Prepared by:	Dillon Alegre, Grant Brink & Rachel Ellison, Environmental Assessment Services, LLC
Report Period:	August 1 st to August 15 th , 2023
Re:	CRAMER FISH SCIENCES - WILLAMETTE VALLEY FISH PASSAGE MONITORING VIA ROTARY SCREW TRAPS

Project Schedule

Site	Task	Start	End	Days
Breitenbush River RST	Trap Install	6/16/2023	6/16/2023	1
Breitenbush River RST	Operation	6/16/2023	11/30/2023	167
Breitenbush River RST	Trapping Efficiency (749 fish)	6/21/2023	6/21/2023	1
Breitenbush River RST	Trapping Efficiency (763 fish)	7/6/2023	7/6/2023	1
Breitenbush River RST	Trapping Efficiency (791 fish)	8/2/2023	8/2/2023	1
Detroit Head of Reservoir- North Santiam River RST	Trap Install	4/19/2023	4/19/2023	1
Detroit Head of Reservoir- North Santiam River RST	Operation	5/4/2023	11/30/2023	210
Detroit Head of Reservoir- North Santiam River RST	Trapping Efficiency (539 fish)	6/6/2023	6/6/2023	1
Detroit Head of Reservoir- North Santiam River RST	Trapping Efficiency (750 fish)	6/20/2023	6/20/2023	1
Detroit Head of Reservoir- North Santiam River RST	Trapping Efficiency (750 fish)	7/6/2023	7/6/2023	1
Detroit Head of Reservoir- North Santiam River RST	Trapping Efficiency (750 fish)	8/2/2023	8/2/2023	1
Green Peter Head of Reservoir- Middle Santiam River RST	Highline Install	4/25/2023	4/25/2023	1
Green Peter Head of Reservoir- Middle Santiam River RST	Trap Install	4/26/2023	4/26/2023	1
Green Peter Head of Reservoir- Middle Santiam River RST	Operation	5/4/2023	11/30/2023	210
Green Peter Head of Reservoir- Middle Santiam River RST	Trapping Efficiency	6/7/2023	6/7/2023	1

Table 1. Project Schedule

	(1000 dead,			
	750 alive)			
	Trapping			
Green Peter Head of Reservoir- Middle Santiam River RST	Efficiency	7/28/2023	7/28/2023	1
	(750)			
	Trapping			
Green Peter Head of Reservoir- Middle Santiam River RST	Efficiency	8/16/2023	8/16/2023	1
	(1,008 live fish)			
Lookout Dam Tailrace RSTs	Operation	8/01/2023	12/31/2023	152
Hills Creek Head of Reservoir RST	Trap Install	5/9/2023	5/9/2023	1
Hills Creek Head of Reservoir RST	Operation	5/9/2023	6/30/2023	52
Hills Creek Head of Reservoir RST	Removal	6/30/2023	6/30/2023	1
	Trapping			
Hills Creek Head of Reservoir RST	Efficiency	5/18/2023	5/18/2023	1
	(519 fish)			
	Trapping			
Hills Creek Head of Reservoir RST	Efficiency	6/19/2023	6/19/2023	1
	(760 fish)			

Table 2. Sampling Dates for Reporting Period

		-	_	_	
	Total	Current	Current	Days	Total
Site	Sampling	Reporting	Reporting	Sampled This	Days
	Period Start	Period Start	Period End	Period	Sampled
Breitenbush River RST	6/16/2023	8/1/2023	8/15/2023	15	76
Detroit Head of Reservoir- North					
Santiam River RST	5/4/2023	8/1/2023	8/15/2023	15	102
Green Peter Head of Reservoir-	5/4/2023	8/1/2023	8/15/2023	15	102
Middle Santiam River RST	5/4/2025	0/1/2023	0/15/2025	15	102
Lookout Point Dam PH	8/1/2023	8/1/2023	8/15/2023	40	10
Lookoul Point Dam PH	8/1/2023	8/1/2023	8/15/2023	13	13
Lookout Point Dam Spill	8/1/2023	8/1/2023	8/15/2023	13	13
Hills Creek Head of Reservoir RST	5/9/2023	8/1/2023	8/15/2023	0	52

Site	Species	Catch (Reporting Period)	Recaptures (Reporting Period)	Total Catch
Breitenbush River RST	CHS	8	12	48
Breitenbush River RST	STW	3	0	320
Detroit Head of Reservoir- North Santiam River RST	CHS	29	19	9353
Detroit Head of Reservoir- North Santiam River RST	STW	11	0	559
Green Peter Head of Reservoir- Middle Santiam River RST	CHS	0	0	21
Green Peter Head of Reservoir- Middle Santiam River RST	STW	0	0	0
Lookout Point Dam	CHS	3	0	0
Hills Creek Head of Reservoir RST	CHS	0	0	93

Table 3. Willamette Valley Rotary Screw Trap Monitoring Catch Summary

Summary of Rotary Screw Trap Data

For this reporting period, traps were operated at the following 4 locations: Detroit Head of Reservoir – North Santiam River, Breitenbush River, Green Peter Head of Reservoir – Middle Santiam River, and Lookout Dam Tailrace.

The Detroit Head of Reservoir – North Santiam RST and Green Peter Head of Reservoir – Middle Santiam RST were installed on April 19th and 26th, respectively. The RSTs at Detroit Head of Reservoir – North Santiam and Green Peter Head of Reservoir – Middle Santiam rivers started sampling on May 4th once permits were received. The Hills Creek Head of Reservoir RST on the upper Middle Fork Willamette River was installed and began sampling on May 9th. Sampling concluded at the Hills Creek Head of Reservoir site on June 30th and was removed for the remainder of the year. The RST for the Breitenbush River was installed on June 16th and began sampling on the same day.

The RSTs in the Lookout Dam Tailrace began sampling under contract W9127N19D0009 on August 1, 2023. Sampling at Lookout Dam Tailrace prior to August 1, 2023 was conducted by EAS for the USACE under contract W9127N19D0007. This report reflects research conducted starting August 1st, 2023 but will include season totals from January 1st, 2023 onward.

Winter Steelhead may be present at the Breitenbush River, Detroit Head of Reservoir – North Santiam River, and Green Peter Head of Reservoir – Middle Santiam River sites. All natural origin juvenile *O. mykiss* captured at these sites will be treated and reported as Winter Steelhead.

Sampling start dates are included in Table 2, and season total collection numbers are displayed in Table 3. The locations of the RSTs are depicted in Figures 1 through 5.



Portland Salem Eugene OREGON FIGURE 1 Breitenbush River

RST Locations

•

500 Feet







FIGURE 2 North Santiam Above Detroit



•

N





FIGURE 3 Middle Santiam River



•






FIGURE 4 Lookout Dam Tailrace

EAS ENVIRONMENTAL ASSESSMENT SERVICES Wholly Owned Subsidiary of Natives of Kodiak

500 Feet

Imagery Source: 2019, ESRI.

MAP AREA

OREGON

Portland

Salem

Eugene





NFS-Rd-2120



500 Feet

He Fork Willamette River

NFS-Rd21

Middle Fork

Willamette Above Hills Creek RST

Breitenbush River

The Breitenbush River RST was installed on June 16th, 2023 and began sampling the same day. All natural origin *O. mykiss* captured at this site will be reported as Winter Steelhead.

Target Species

This reporting period began on August 1st and ended on August 15th. There were a total of 8 Chinook Salmon (CHS) and 3 Winter Steelhead (STW) captured during the 15-day sampling period (Figure 6). Sampling duration was 100% of the reporting period for the RST. Figure 7 shows length frequency data to-date. Table 4 provides life stage, length, and weight data for all Chinook Salmon and Winter Steelhead that have been caught at the Breitenbush River site to-date and for the reporting period.



Figure 6. Chinook and Winter Steelhead Captured per day 08/01/2023 to 08/15/2023 (Breitenbush River)




Figure 7. Length Frequency of Juvenile Chinook Sampled Season To-Date (Breitenbush River)

	To-Date (Since June 16, 2023)									
Site	Route	Species	Life	Collected	L	ength (m	m)*		Weight	(g)*
			stage		Min	Мах	Mean	Min	Мах	Mean
		CHS	Fry	10	44	57	48.8	1.0	1.9	1.2
		CHS	Parr	36	46	92	64.3	1.4	8.6	3.4
Breitenbush	5ft	CHS	Smolt	2	91	96	93.5	8.9	10.2	9.6
River		STW	Fry	311	21	47	30.0	N/A	N/A	N/A
		STW	Parr	9	43	120	76.4	1.2	20.0	7.5
		STW	Smolt	0	N/A	N/A	N/A	N/A	N/A	N/A

Table 4. Descriptive Statistics of Target Species Captured at the Breitenbush River To-Date

*Fish that were missing heads or caudal fins are not included in length and weight calculations.

	August 1-15, 2023									
Site	Route	Species	Life	Collected		ength (m	m)*		Weight (g)*
			stage		Min	Max	Mean	Min	Мах	Mean
		CHS	Fry	0	N/A	N/A	N/A	N/A	N/A	N/A
		CHS	Parr	6	64	92	76.5	3.7	8.6	5.3
Breitenbush	5ft	CHS	Smolt	2	91	96	93.5	8.9	10.2	9.6
River		STW	Fry	1	37	37	37.0	N/A	N/A	N/A
		STW	Parr	2	50	51	50.5	1.1	2.2	1.7
		STW	Smolt	0	N/A	N/A	N/A	N/A	N/A	N/A

*Fish that were missing heads or caudal fins are not included in length and weight calculations.

Trapping Efficiency

On 8/2/2023 791 adipose and right ventral fin clipped fish were released above the trap site to evaluate the trapping efficiency of the 5 ft RST. 12 fish were recaptured for an efficiency of 1.5%

Release #	Recapture #	Capture Efficiency
791	12	1.5% (12/791)

Injuries and Copepod Infection

Partial descaling <20% was observed in 5 of the 8 Chinook captured (62.5%), 0 displayed descaling >20% (0.0%), 2 displayed body injury (25.0%), 0 had eye injuries (0.0%), 0 had copepods present in the branchial cavity (0.0%) and 1 had copepods on fins (12.5%). 0 Chinook displayed gas bubble disease (0.0%). There was 0 mortalities (0.0%).

Partial descaling <20% was observed on 0 of the 3 Winter Steelhead captured (0.0%) and 0 displayed descaling >20% (0.0%), 0 displayed body injury (0.0%), 0 had eye injury (0.0%), 0 had copepods present in the branchial cavity (0.0%) and 0 had copepods on fins (0.0%). 0 Winter Steelhead displayed gas bubble disease (0.0%). There were 0 mortalities (0.0%). Injury data is summarized in table 5.

Table 5. Number of Descaled, Bodily/Eye Injured, Copepod Infected and Dead ChinookSalmon and Winter Steelhead for Sampling Period (Breitenbush River).

Site	Species	# Fish Collected	# DSC* <20%	# DSC* >20%	# with Body Injuries	# with Eye Injuries	# with COP* In B.C.	# with COP* on Fins	Mortalities
Breitenbush	Chinook	8	5	0	2	0	0	1	0
River	Winter Steelhead	3	0	0	0	0	0	0	0

*DSC=Descaled, COP=Copepods, B.C.=Branchial Cavity

Collected DNA and Scale Samples

DNA was collected from 8 Spring Chinook and 3 Winter Steelhead. Scale samples were collected from 8 Spring Chinook and 3 Winter Steelhead.

PIT Tags

7 fish were PIT tagged during this reporting period. More information regarding PIT tagged fish can be found in Appendix D.

VIE Marking

Visible Implant Elastomer (VIE) trials commenced on 6/16/2023. VIE tag color is changed every month to distinctly mark groups of fish by capture date. Since then, 28 Spring Chinook and 10 Winter Steelhead have been marked with fluorescent elastomer. No fish have been recaptured at downstream sites to date.

Date Tagged	Species	Tag Location	VIE Color	# Tagged	# Recaptured to Date
6/16/2023-6/30/2023	Chinook	Head	Pink	23	0
7/1/2023-7/15/2023	Chinook	Head	Green	2	0
7/16/2023-7/31/2023	Chinook	Head	Green	2	0
7/16/2023-7/31/2023	O. mykiss	Head	Green	7	0
8/1/2023-8/15/2023	Chinook	Head	Yellow (2x)	1	0
8/1/2023-8/15/2023	O. mykiss	Head	Yellow (2x)	3	0

Non-Target Species

3 non-target species were captured during this reporting period. A summary of non-target fish capture is provided in table 6.

Species	5 ft Capture	5 ft Mortality	Season Total	Season Total Mortality				
Kokanee	0	0	0	0				
Cutthroat Trout	0	0	0	0				
O. mykiss (clipped)	1	0	9	5				
Sculpin	2	0	9	2				
Totals	3	0	15	7				

Table 6. Summary of Non-target Species (Breitenbush River).

Stream Statistics


Basic stream statistics at the proposed Breitenbush River RST site were calculated from data downloaded from the U.S. Geological Survey stream gauge number 14179000. Instantaneous discharge (cfs) and Gauge height (feet) flow metrics are available at this gauge. During the reporting period, daily maximum values for instantaneous discharge ranged from 116.0 cfs to 133.0 (mean: 125.3 cfs). Figure 8 shows instantaneous discharge.

Stream temperatures will be recorded every 2 hours for the length of the reporting period for the RST (Figure 9).


Catch per unit of effort (CPUE) data are summarized in Table 7. Gage height and capture data for the duration of monitoring efforts at this location are provided in Appendix B.

	Chinook	Winter Steelhead
Description	(5 ft)	(5 ft)
Catch	8	3
Effort (hrs)	356.9	356.9
CPUE (fish/hr)	0.017	0.008

Table 7. Summary of salmonid CPUE, Breitenbush River.









North Santiam River – Detroit Head of Reservoir

The Detroit Head of Reservoir- North Santiam River RST was installed on April 19th, 2023. This site started sampling on May 4, 2023. All natural origin *O. mykiss* captured at this site will be reported as Winter Steelhead.

Target Species

This reporting period began on August 1st and ended on August 15th. There were a total of 29 Chinook Salmon (CHS) and 11 Winter Steelhead (STW) captured during the 15-day sampling period (Figure 10). Sampling duration was 100% of the reporting period for the RST. Figure 11 shows length frequency data to-date. Table 8 provides life stage, length, and weight data for all Chinook Salmon and Winter Steelhead that have been caught at the Detroit Head of Reservoir site to-date and for the reporting period.




Figure 10. Chinook and Winter Steelhead Captured per day 08/01/2023 to 08/15/2023 (Detroit Head of Reservoir)




Figure 11. Length Frequency of Juvenile Chinook and Winter Steelhead Sampled Season To-Date (Detroit Head of Reservoir)

Table 8. Descriptive Statistics of Target Species Captured at Detroit Head of Reservoir Season To-Date

	To-Date (Since May 04, 2023)											
Site Ro	Route	Species	es Life	Collected	L	Length (mm)*			Weight (g)*			
			stage		Min	Max	Mean	Min	Мах	Mean		
		CHS	Fry	9117	28	60	35.5	N/A	N/A	N/A		
		CHS	Parr	233	41	87	53.2	1.0	7.6	1.9		
Detroit	5ft	CHS	Smolt	3	61	86	75.0	2.4	6.1	4.5		
HOR		STW	Fry	544	17	49	34.3	N/A	N/A	N/A		
		STW	Parr	10	45	99	69.8	1.0	10.6	5.0		
		STW	Smolt	2	188	408	298	66.5	66.5	66.5		

*Fish that were missing heads are not included in length and weight calculations.

	August 1-15, 2023											
Site R	Route	Specie	Life	Collected	L	Length (mm) [*]			Weight (g) [*]			
			s stage		Min	Мах	Mean	Min	Мах	Mean		
	CHS	Fry	1	48	48	48.0	1.3	1.3	1.3			
		CHS	Parr	26	46	72	55.8	1.0	3.7	2.0		
Detroit HOR	5ft	CHS	Smolt	2	61	86	73.5	2.4	6.1	4.3		
nok		STW	Fry	7	23	48	33.9	<1	1.4	N/A		
		STW	Parr	4	45	52	47.8	1.0	1.5	1.2		
		STW	Smolt	0	N/A	N/A	N/A	N/A	N/A	N/A		

*Fish that were missing heads are not included in length and weight calculations.

Trapping Efficiency

On 8/2/2023 750 adipose and left ventral fin clipped fish were released above the trap site to evaluate the trapping efficiency of the 5 ft RST. 19 fish were recaptured for an efficiency of 2.5%

Detroit Head of Reservoir	Release #	Recapture #	Capture Efficiency
5 0 T	750	40	2.5%
5ft Trap	750	19	(19/750)

Injuries and Copepod Infection

Partial descaling <20% was observed in 8 of the 29 Chinook captured (27.6%), 0 displayed descaling >20% (0.0%), 6 displayed body injury (20.7%), 0 had eye injuries (0.0%), 0 had copepods present in the branchial cavity (0.0%) and 0 had copepods on fins (0.0%). 0 Chinook displayed gas bubble disease (0.0%). There were 0 mortalities (0.0%).

Partial descaling <20% was observed on 0 of the 11 Winter Steelhead captured (0.0%) and 0 displayed descaling >20% (0.0%), 1 displayed body injury (9.1%), 0 had eye injuries (0.0%), 0 had copepods present in the branchial cavity (0.0%) and 0 had copepods on fins (0.0%). 0 Winter Steelhead displayed gas bubble disease (0.0%). There were 0 mortalities (0.0%). Injury data is summarized in table 9.

Table 9. Number of Descaled, Bodily/Eye Injured, Copepod Infected and Dead ChinookSalmon and Winter Steelhead for Sampling Period (Detroit Head of Reservoir).

Site	Species	# Fish Collected	# DSC* <20%	# DSC* >20%	# with Body Injuries	# with Eye Injuries	# with COP* In B.C.	# with COP* on Fins	Mortalities
Detroit	Chinook	29	8	0	6	0	0	0	0
HOR	Winter Steelhead	11	0	0	1	0	0	0	0

*DSC=Descaled, COP=Copepods, B.C.=Branchial Cavity

Collected DNA and Scale Samples

For the reporting period, DNA was collected from 29 Spring Chinook and 5 Winter Steelhead. Scale samples were collected from 25 Spring Chinook and 1 Winter Steelhead. The other targets captured did not meet length criteria for DNA sampling or were too descaled/damaged to collect samples.

PIT Tags

4 Spring Chinook and 0 Winter Steelhead were PIT tagged during this reporting period. More information regarding PIT tagged fish can be found in Appendix D.

VIE Marking

Visible Implant Elastomer (VIE) trials commenced at the Detroit Head of Reservoir – North Santiam River site on 5/5/2023. VIE tag color is changed every month to distinctly mark groups of fish by capture date. Since then, 5,385 Chinook and 326 Winter Steelhead have been VIE marked with fluorescent elastomer. No fish with VIE marks have been detected at downstream RST sites to date. Fish still showing an egg sac are not VIE marked.

Date Tagged	Species	Tag Location	VIE Color	# Tagged	# Recaptured to Date
5/01/2023-5/15/2023	Chinook	Right Dorsal	Orange	889	0
5/01/2023-5/15/2023	O. mykiss	Right Dorsal	Orange	60	0
5/16/2023- 5/31/2023	Chinook	Right Dorsal	Orange	2,700	0
5/16/2023- 5/31/2023	O. mykiss	Right Dorsal	Orange	237	0
6/1/2023-6/15/2023	Chinook	Right Dorsal	Pink	1048	0
6/1/2023-6/15/2023	O. mykiss	Right Dorsal	Pink	21	0
6/16/2023-6/30/2023	Chinook	Right Dorsal	Pink	539	0
7/1/2023-7/15/2023	Chinook	Right Dorsal	Green	110	0
7/16/2023-7/31/2023	Chinook	Right Dorsal	Green	74	0
7/16/2023-7/31/2023	O. mykiss	Right Dorsal	Green	1	0
8/1/2023-8/15/2023	Chinook	Right Dorsal	Yellow (2x)	25	0
8/1/2023-8/15/2023	O. mykiss	Right Dorsal	Yellow (2x)	7	0

Non-Target Species

5 non-target species fish were captured during the reporting period; the data is summarized below in table 10.

Species	5 ft Capture	5 ft Mortality	Season Total	Season Total Mortality
Kokanee	0	0	81	1
Chinook (clipped)	2	0	4	0
Cutthroat Trout	0	0	1	0
Sculpin	1	0	13	2
Mountain Whitefish	0	0	3	0
O. mykiss (clipped)	0	0	6	0
Dace	1	0	3	0
Unknown	1	0	2	1
Totals	5	0	113	4

Table 10. Summary of Non-target Species (Detroit Head of Reservoir).

Stream Statistics

Basic stream statistics at the Detroit Head of Reservoir site were calculated from data downloaded from U.S. Geological Survey stream gauge number 14178000. Gauge height (feet) and Discharge (cfs) metrics are provided at gauge 14178000. During the reporting period, daily maximum values for instantaneous discharge ranged from 426.0 cfs to 456.0 cfs (mean: 436.0 cfs) during the reporting period. Figure 12 shows instantaneous discharge.

Stream temperatures were recorded every 2 hours for the length of the reporting period at the Detroit Head of Reservoir RST site. Figure 13 shows temperature during the reporting period.

Catch per unit of effort (CPUE) data are summarized in Table 11. Discharge and capture data for the duration of monitoring efforts at this location are provided in Appendix B.

	Chinook	Winter Steelhead							
Description	(5 ft)	(5 ft)							
Catch	29	12							
Effort (hrs)	357.7	357.7							
CPUE (fish/hr)	0.081	0.034							

Table 11. Summary of salmonid CPUE, Detroit Head of Reservoir – North Santiam River.




Figure 12. Discharge (cfs); Detroit Head of Reservoir – North Santiam River




Figure 13. Temperature at RST (Detroit Head of Reservoir – North Santiam River)

Middle Santiam River- Green Peter Head of Reservoir

The Green Peter Head of Reservoir- Middle Santiam River RST was installed on April 26th, 2023. This site started sampling on May 4th, 2023. All natural origin *O. mykiss* captured at this site will be reported as Winter Steelhead.

Target Species

This reporting period began on August 1st and ended on August 15th. There were a total of 0 Chinook Salmon (CHS) and 0 Winter Steelhead (STW) captured during the 15-day sampling period (Figure 14). Sampling duration was 100% of the reporting period for the RST. Figure 15 shows length frequency data to-date. Table 12 provides life stage, length, and weight data for all Chinook Salmon and Winter Steelhead that have been caught at the Middle Santiam River- Green Peter Head of Reservoir site to-date and for the reporting period.




Figure 14. Chinook Captured per day 08/01/2023 to 08/15/2023 (Green Peter Head of Reservoir – Middle Santiam River)




Figure 15. Length Frequency of Juvenile Chinook Sampled Season To-Date (Green Peter Head of Reservoir – Middle Santiam River)

	To-date (since May 04, 2023)											
Site	Route	Species	Life	Collected	L	.ength (m	m)*	Weight (g)*				
			stage		Min	Мах	Mean	Min	Мах	Mean		
		CHS	Fry	21	33	45	36.4	N/A	N/A	N/A		
Green	5ft	CHS	Parr	0	N/A	N/A	N/A	N/A	N/A	N/A		
Peter Head of		CHS	Smolt	0	N/A	N/A	N/A	N/A	N/A	N/A		
Reservoir		STW	Fry	1	36	36	36	N/A	N/A	N/A		
-Middle Santiam		STW	Parr	0	N/A	N/A	N/A	N/A	N/A	N/A		
		STW	Smolt	0	N/A	N/A	N/A	N/A	N/A	N/A		

Table 12. Descriptive Statistics of Target Species Captured at Green Peter Head of Reservoir – Middle Santiam River Season To-Date

*Fish that were missing heads are not included in length and weight calculations.

	August 1-15, 2023												
Site	Route	Route Species		Collected	L	.ength (m	m)*	Weight (g) [*]					
			stage		Min	Мах	Mean	Min	Мах	Mean			
		CHS	Fry	0	N/A	N/A	N/A	N/A	N/A	N/A			
Green		CHS	Parr	0	N/A	N/A	N/A	N/A	N/A	N/A			
Peter Head of	5ft	CHS	Smolt	0	N/A	N/A	N/A	N/A	N/A	N/A			
Reservoir -Middle		STW	Fry	0	N/A	N/A	N/A	N/A	N/A	N/A			
Santiam		STW	Parr	0	N/A	N/A	N/A	N/A	N/A	N/A			
		STW	Smolt	0	N/A	N/A	N/A	N/A	N/A	N/A			

*Fish that were missing heads are not included in length and weight calculations.

Trapping Efficiency

On 7/28/2023 750 adipose and left ventral clipped fish were released above the trap site to evaluate the trapping efficiency of the 5 ft RST. 0 fish was recaptured for an efficiency of 0.0%.

Green Peter Head of Reservoir- Middle Santiam River	Release #	Recapture #	Capture Efficiency
5ft Trap	Alive (750)	0	0/750 0.0%
	Dead (0)	N/A	N/A

Injuries and Copepod Infection

Partial descaling <20% was observed in 0 of the 0 Chinook captured (0.0%), 0 displayed descaling >20% (0.0%), 0 displayed body injury (0.0%), 0 had eye injuries (0.0%), 0 had copepods present in the branchial cavity (0.0%) and 0 had copepods on fins (0.0%). 0 Chinook displayed gas bubble disease (0.0%). There was 0 mortality (0.0%).

Partial descaling <20% was observed on 0 of the 0 Winter Steelhead captured (0.0%) and 0 displayed descaling >20% (0.0%), 0 displayed body injury (0.0%), 0 had eye injuries (0.0%), 0 had copepods present in the branchial cavity (0.0%) and 0 had copepods on fins (0.0%). 0 Winter Steelhead displayed gas bubble disease (0.0%). There were 0 mortalities (0.0%). Injury data is summarized in Table 13.

Table 13. Number of Descaled, Bodily/Eye Injured, Copepod Infected and Dead Chinook Salmon and Winter Steelhead for Sampling Period (Green Peter Head of Reservoir-Middle Santiam River).

Site	Species	# Fish Collected	# DSC* <20%	# DSC* >20%	# with Body Injuries	# with Eye Injuries	# with COP* In B.C.	# with COP* on Fins	Mortalities
Green Peter Head of	Chinook	0	0	0	0	0	0	0	0
Reservoir- Middle Santiam	Winter Steelhead	0	0	0	0	0	0	0	0

*DSC=Descaled, COP=Copepods, B.C.=Branchial Cavity

Collected DNA and Scale Samples

For the reporting period, DNA was collected from 0 Spring Chinook and 0 Winter Steelhead. Scale samples were collected from 0 Spring Chinook and 0 Winter Steelhead. The other targets captured did not meet length criteria for DNA sampling or were too descaled/damaged to collect samples.

PIT Tags

0 Spring Chinook and 0 Winter Steelhead were PIT tagged during this reporting period. All fish captured did not meet the size criteria for PIT tagging. More information regarding PIT tagged fish can be found in Appendix D.

VIE Marking

Visible Implant Elastomer (VIE) trials commenced at the Green Peter Head of Reservoir – Middle Santiam River site on 5/5/2023. VIE tag color and locations are changed every month to distinctly mark groups of fish by capture date. Since then, 15 Chinook and 1 Winter Steelhead have been VIE marked with fluorescent elastomer. No fish with VIE marks have been detected at downstream RST sites to date. Fish still showing an egg sac are not VIE marked.

Date Tagged	Species	Tag Location	VIE Color	# Tagged	# Recaptured to Date
5/01/2023-5/15/2023	Chinook	Right Dorsal	Orange	14	0
5/01/2023-5/15/2023	O. mykiss	Right Dorsal	Orange	1	0
5/16/2023-5/31/2023	Chinook	Right Dorsal	Orange	1	0

Non-Target Species

1 non-target fish were collected during the reporting period; the data is summarized below in Table 14.

Table 14. Summary of Non-target Species (Green Peter Head of Reservoir – Middle Santiam River).

Species	5 ft Capture	5 ft Mortality	Season Total	Season Total Mortality
Kokanee	0	0	5	0
Cutthroat Trout	0	0	0	0
Dace	1	0	17	0
Sculpin	0	0	9	0
Totals	1	0	31	0

Stream Statistics

Basic stream statistics at the Green Peter Head of Reservoir – Middle Santiam River site were calculated from data downloaded from the U.S. Geological Survey stream gauge number 14185800. Gauge height

(feet) is the only flow metric available at this gauge. During the reporting period, daily maximum values for gage height ranged from 0.9 ft to 1.0 ft (mean: 1.0 ft). Figure 16 shows gage height.

Stream temperatures were recorded every 2 hours for the length of the report period for the RST (Figure 17). Temperature probes for the trap operated normally throughout this reporting period.

Catch per unit of effort (CPUE) data are summarized in Table 15. Gage height and capture data for the duration of monitoring efforts at this location are provided in Appendix B.

Table 15. Summary of salmonid CPUE, Green Peter Head of Reservoir – Middle Santiam River.

	Chinook	Winter Steelhead
Description	(5 ft)	(5 ft)
Catch	0	0
Effort (hrs)	358.6	358.6
CPUE (fish/hr)	0	0




Figure 16. Gage Height (feet); Green Peter Head of Reservoir - Middle Santiam River




Figure 17. Temperature at RST (Green Peter Head of Reservoir – Middle Santiam River)

Middle Fork Willamette – Lookout Dam Tailrace

The RSTs in the Lookout Dam Tailrace began sampling under contract W9127N19D0009 on August 1, 2023. Sampling at Lookout Dam Tailrace prior to August 1, 2023 was conducted by EAS for the USACE under contract W9127N19D0007. This report reflects research conducted starting August 1st, 2023 but will include season totals from January 1st, 2023 onward.

Target Species

The reporting period began August 1st and ended on August 15th. 3 Chinook salmon were captured during the 15-day sampling period (Figure 18). The powerhouse traps and the spill trap were raised to the non-sampling position on August 10th due to temperature thresholds. They were lowered the next day on August 11th. They were raised again on August 15th due to air quality safety concerns from the Bedrock and Lookout fires. The traps were operated 86.7% of the reporting period. Table 16 provides life stage, length, and weight data for all Chinook salmon that have been caught at the Lookout Point Dam Tailrace site to-date and Figure 19 shows length frequency data to-date.




Figure 18. Chinook Captured Per Day 08/01/2023 to 08/15/2023 (Lookout Point Dam Tailrace).




Figure 19. Length Frequency of Juvenile Chinook Sampled in 2023 (Lookout Point Dam Tailrace).

To-Date (Since Jan. 1, 2023)										
	_				Length (mm)*			V	Veight (g)*
Site	Route	Species	Life stage	Collected	Min	Max	Mean	Min	Мах	Mean
	PH 1	CHS	Smolt	14	109	275	167.6	14.8	269.0	66.6
		CHS	Parr	1	96	96	96	9.4	9.4	9.4
		CHS	Fry	1	52	52	52	3.0	3.0	3.0
		CHS	Smolt	4	113	250	160.3	16.5	194.6	69.9
Lookout Point Dam	PH 2	CHS	Parr	4	57	86	71.0	1.8	7.6	5.3
		CHS	Fry	4	33	37	34.8	N/A	N/A	N/A
		CHS	Smolt	21	122	247	172.2	19.7	161.4	60.7
	Spill	CHS	Parr	1	125	125	125	17.6	17.6	17.6
		CHS	Fry	2	44	55	49.5	1.0	1.6	1.3
				August 1-15	, 2023	1				
Site	Route	Species	Life	Collected	Length (mm)*			Weight (g) [*]		
U IIO	nouto	opooloo	stage		Min	Мах	Mean	Min	Мах	Mean
	PH 1	CHS	Smolt	1	170	170	170.0	65.5	65.5	65.5
		CHS	Parr	0	N/A	N/A	N/A	N/A	N/A	N/A
		CHS	Fry	0	N/A	N/A	N/A	N/A	N/A	N/A
		CHS	Smolt	0	N/A	N/A	N/A	N/A	N/A	N/A
Lookout Point Dam	PH 2	CHS	Parr	0	N/A	N/A	N/A	N/A	N/A	N/A
		CHS	Fry	0	N/A	N/A	N/A	N/A	N/A	N/A
		CHS	Smolt	2	155	155	155.0	45.2	45.2	45.2
	Spill	CHS	Parr	0	N/A	N/A	N/A	N/A	N/A	N/A
		CHS	Fry	0	N/A	N/A	N/A	N/A	N/A	N/A

Table 16. Descriptive Statistics of Target Species Captured at Lookout Point Dam Tailrace, Season To-Date and for the Reporting Period.

*Some fry are too small to accurately weigh and are omitted from the above tables. **Season totals include sampling completed on the RST project in 2023.

Trapping Efficiency

A total of 4,012 juvenile hatchery Chinook (parr) were adipose clipped and released on 7/18/2023 below Lookout Point Dam. Fish were released in small groups directly into powerhouse flow. A total of 1 fish was recaptured in the traps for an efficiency of 0.02%. Trap specific efficiencies are as follows: 0

recaptured at the PH 1 RST for an efficiency of 0%, 0 recaptured at PH 2 for an efficiency of 0%, and 1 recaptured at the Spill RST for an efficiency of 0.02%.

Lookout Dam (7/18/2023)	Release #	Recapture #	Capture Efficiency
Powerhouse	4,012	1	0.02% (1/4,012)

24-Hour Post Collection Holding Trial

0 Spring Chinook was captured during the current reporting period and held for 24 hours. 0 fish was held from the PWR RST and 0 fish were held from the Spill RST. 0 hold fish died from the PWR RSTs (0.0%). 0 of the fish from Spill RST died during holding (0.0%).

The 1 live target fish was released after a visual assessment due to temperature thresholds being reached.

Injuries and Copepod Infection

There were 2 Chinook captured in the Spill Channel RST. Partial descaling <20% was observed on 0 of 2 Chinook collected at the Spill RST (0.0%), and descaling >20% was observed on 2 of the Chinook collected (100.0%). 2 displayed body injuries (100.0%) and 0 had eye injuries (0.0%). 0 of the Spill RST Chinook had copepods present in the branchial cavity (0.0%) and 0 had copepods present on fins (0.0%). 0 of the fish captured in the Spill RST displayed Gas Bubble Disease (0.0%).

There was 1 Chinook captured in the Powerhouse channel RSTs. Partial descaling <20% was observed on 0 of the 1 Chinook collected at the PWR RSTs (0.0%). Descaling >20% was observed on 1 of the Chinook collected (100.0%). 1 PWR RST fish had bodily injury (100.0%) and 0 had eye injuries (0.0%). 1 of the fish had copepods present in the branchial cavity (100.0%) and 0 had copepods present on fins (0.0%). 0 fish displayed Gas Bubble Disease (0.0%).

There was 1 chinook mortality collected in the Spill RST (50.0%) and 1 in the PWR RSTs (100.0%). Injuries are displayed in Table 17. To date injury data can be found in Appendix A.

Site	Route	# CHS Collected	# DSC* <20%	# DSC* >20%	# with Body Injuries	# with Eye Injuries	# with COP* In B.C.	# with COP* on Fins	Mortalities
Lookout Point Dam	PWR	1	0	1	1	0	1	0	1
Tailrace	Spill	2	0	2	2	0	0	0	1

Table 17. Number of Descaled, Bodily/Eye Injured, Copepod Infected and Dead Chinook Salmon for Sampling Period (Lookout Point Dam Tailrace).

*DSC=Descaled, COP=Copepods, B.C.=Branchial Cavity

Collected DNA and Scale Samples

DNA was collected from 2 Spring Chinook for the reporting period. Scales were collected from 1 Spring Chinook. The other targets captured only received a visual inspection due to water temperature exceeding temperature thresholds.

PIT Tags

No Spring Chinook were PIT tagged during this reporting period. The first 60 target fish per week are prioritized for the 24-Hour Post Collection Holding Study. These fish are not tagged to not bias the results of the holding study. More information regarding PIT tagged fish can be found in Appendix D.

VIE Marking

No VIE marked Spring Chinook have been detected at this site to date.

Non-Target Species

6,111 non-target species were captured during the reporting period; the data is summarized below in Table 18.

Species	PWR Capture	PWR Mortality	Spill Capture	Spill Mortality	Season Total*	Season Total Mortality*
Bass Unknown	5560	5540	53	29	23438	22348
Bluegill	0	0	0	0	59	11
Brown Bullhead	0	0	0	0	3	1
Chinook (clipped)	0	0	0	0	14	2
Crappie	263	97	203	47	155119	109801
Cutthroat Trout	0	0	0	0	0	0
Dace	0	0	0	0	0	0
Largemouth Bass	0	0	0	0	23	23
Largescale Sucker	0	0	0	0	6	4
Northern Pikeminnow	0	0	0	0	4	3
O. mykiss	0	0	1	1	7	1
O. mykiss (clipped)	0	0	0	0	1	1
Pumpkinseed	0	0	0	0	1	0
Redside Shiner	0	0	0	0	1	0
Sculpin	1	0	1	1	135	9
Smallmouth Bass	24	24	3	2	141	133
Spotted Bass	0	0	0	0	1	0
Unknown	0	0	0	0	7	0
Walleye	1	0	1	1	37	8
Totals	5849	5661	262	81	178997	132345

Table 18. Summary of Non-target Species (Lookout Point Dam Tailrace).

*Season totals include sampling completed on the RST project in 2023.

Stream Statistics

Basic stream statistics at Lookout Dam Tailrace site were calculated from data downloaded from the U.S. Geological Survey stream gauge number 14149010. Total dissolved gas saturation or dissolved oxygen concentration measurements are not available at this stream gauge site, or any nearby stream gauges. Gauge height (feet) is the only metric provided at this gauge. During the reporting period, daily maximum values for instantaneous gauge height ranged from 693.2 feet to 693.5 feet (mean: 693.3 feet). Figure 20 shows instantaneous gauge height.

Stream temperatures were recorded every 2 hours using temperature probes at the PWR and Spill Lookout Dam RST's during this reporting period. Temperature probes operated normally, and the data is shown below in (Figure 21 and Figure 22).

Flows through the Powerhouse and Spill during the reporting period averaged 801.4 and 1,215.2 cubic feet per second (cfs) respectively (Figure 23). Catch per unit of effort (CPUE) data are summarized in Table 19. Discharge and capture data for the duration of monitoring efforts at this location are provided in Appendix B.

	Chinook					
Description	PH 1	PH 2	Spill			
Catch	1	0	2			
Effort (hrs)	333.8	335.7	333.5			
CPUE (fish/hr)	0.003	0	0.006			

 Table 19. Summary of Chinook CPUE at Lookout Point Dam Tailrace.




Figure 20. Gauge Height (feet); below Lookout Dam.




Figure 21. Temperature at RST (Lookout Dam PWR).




Figure 22. Temperature at RST (Lookout Dam Spill).




Figure 23. Hourly Flows PWR vs. Spill (Lookout Dam Tailrace)

Middle Fork Willamette River-Hills Creek Head of Reservoir

The Hills Creek Head of Reservoir RST was installed and began sampling on May 9th, 2023. Sampling concluded at this site on June 30, 2023 and the RST was removed.

Target Species

A total of 93 Spring Chinook were captured during sampling in 2023. Figure 24 shows length frequency data of captured Chinook for sampling in 2023. Table 20 provides life stage, length, and weight data for all Chinook Salmon that have been caught at the Middle Fork Willamette River- Hills Creek Head of Reservoir site to-date and for the reporting period.




Figure 24. Length Frequency of Juvenile Chinook Sampled Season To-Date (Hills Creek Head of Reservoir)

Table 20. Descriptive Statistics of Target Species Captured at Hills Creek Head ofReservoir Season To-Date

			To-D	ate (Since M	ay 09,	2023)				
Site	Route	Species	Life	Collected	Le	ength (n	nm) [,]		Weight	(g) [.]
			stage		Min	Мах	Mean	Min	Мах	Mean
Hills Creek		CHS	Fry	60	30	50	38.9	<1	2.5	1.4
Head of Reservoir	5 ft	CHS	Parr	33	38	76	52.6	1.0	6.0	2.1
		CHS	Smolt	0	N/A	N/A	N/A	N/A	N/A	N/A

*Most fry are too small to collect accurate weights and thus some metrics are not available for them.

Trapping Efficiency

On May 18th, 519 adipose clipped and PIT Tagged fish were released for a trapping efficiency trial at the Hills Creek Head of Reservoir site. 44 fish were recaptured in the RST for a trapping efficiency of 8.5%

Hills Creek Head of Reservoir	Release #	Recapture #	Capture Efficiency
5# Teen	540		8.5%
5ft Trap	519	44	(44/519)

On June 19th, 760 adipose clipped and PIT Tagged fish were released for a trapping efficiency trial at the Hills Creek Head of Reservoir site. 6 fish were recaptured in the RST for a trapping efficiency of 0.79%.

Hills Creek Head of Reservoir	Release #	Recapture #	Capture Efficiency
5ft Trap	760	6	0.8% (6/760)

PIT Tags and VIE Marking

A total of 3 fish were PIT tagged and 71 fish were VIE marked at the Hills Creek Head of Reservoir-Middle Fork Willamette site in 2023. No tagged or VIE marked fish were redetected downstream. Table 21 provides a summary of VIE marked fish at the Hills Creek Head of Reservoir- Middle Fork Willamette River site.

Table 21. Summary of VIE marked Chinook at the Hills Creek Head of Reservoir- MiddleFork Willamette site in 2023.

Date Tagged	Species	Tag Location	VIE Color	# Tagged	# Recaptured to Date
5/1/2023-5/30/2023	Chinook	Left Dorsal	Orange	19	0
5/1/2023-5/30/2023	Chinook	Right Dorsal	Orange	11	0
6/1/2023-6/30/2023	Chinook	Left Dorsal	Pink	37	0
6/1/2023-6/30/2023	Chinook	Right Dorsal	Pink	4	0

Non-Target Species

A total of 232 non-target species fish were captured during sampling in 2023; the data is summarized below in Table 22.

Table 22. Summary of Non-target Species (Hills Creek Head of Reservoir).

Species	Season Total	Season Total Mortality
Dace	87	1
Cutthroat Trout	2	0
O. mykiss	26	0
Bull Trout	1	0
Brook Lamprey	18	2
Sculpin	20	1
Largescale Sucker	64	1
Mountain Whitefish	2	0
Redside Shiner	12	0
Totals	232	5

Issues Encountered

The Lookout Dam Tailrace traps were raised to the non-sampling position on August 10th due to temperature thresholds. The Lookout Dam Tailrace traps were also raised to the non-sampling position on August 15th due to air quality above our safety thresholds. The Lowell to Oakridge corridor is currently reading over 400 on the AQI scale.

Upcoming USACE Support Services

None at this time.

Appendix A
Chinook (CHS) To-Date

						C	hinod	ok Inju	ries	to-da	ate											
Site/Trap/Life Stage	Total Fish	MUNK	DS<2	BLO	EYB	FUN	BKD	сор	DS>2	PRD	FID HRO	BO	ЮН	вит	НВР	BRU	TEA	ОРD	HIN	FVB	POP	GBD
Breitenbush River																						
5 ft	48		14			1		2			13					1						
Parr	36		12			1		2			8					1						
Smolt	2		2								2											
Fry	10										3											
Detroit HOR																	30	38				
5 ft	9354	1	62		11	1			8	9	78		1	17		57	30	38	25	43	9	
Parr	233		33			1					28					2	3					
Smolt	3		2								1											
Fry	9118	1	27		11				8	9	49		1	17		55	27	38	25	43	9	
Green Peter HOR	21																					
5 ft	21										1								1			
Fry	21										1								1			
Lookout Dam Tail.	131		62		21			33	51		86					14		22	19	23		16
PH 1	44		18	1	8	1	1	15	20		32			5	2	9	4	9	7	5		5
Parr	4		1		1				2		2				1	1		1				
Smolt	39		17	1	7	1	1	15	18		30			5	1	8	4	8	7	5		5
Fry	1																					
PH 2	25		14		6		1	3	7		16			1		2	3	2	5	3		
Parr	8		6		4				2		6					1	1	1	3			
Smolt	13		8		2		1	3	4		10			1		1		1	1	2		
Fry	4								1								2		1	1		
Spill	62		30		7	1	3	15	24		38			1		3	2	11	7	15	3	11
Parr	7		2						3		2											
Smolt	53		27		6	1	3	15	21		35			1		3	2	11	7	15	3	11
Fry	2		1		1						1											
Hills Creek HOR	93																					
5 ft	93		6							1	2											
Parr	33		4							1	2											
Fry	60		2																			

Chinook (CHS)	During	Reporting Period	
---------------	--------	-------------------------	--

	C	hinook Inju	ries During Re	eporting Peri	od 8-1-2023 to	o 8-15-2023		
Site/Trap/Life Stage	Total Fish ≥	DS<2 BLO	EYB FUN BKD	COP DS>2 PRD	FID HBO	BU BVT HBP	BRU TEA OPD	FVB POP GBD
Breitenbush River								
5 ft	8	6		1	3			
Parr	6	4		1	1			
Smolt	2	2			2			
Detroit HOR								
5 ft	29	8			6			
Fry	1							
Parr	26	7			5			
Smolt	2	1			1			
Lookout Dam Tailrace								2
PH 1	3			1 3	3	2	1 1 2	2
Smolt	1			1 1	1	1	1 1	1
Spill	2			2	2	1	2	1
Smolt	2			2	2	1	2	1

Steelhead (O. mykiss) To Date


					O. my	kiss Inju	uries	to-d	ate									
Site/Trap/Life Stage	Total Fish	MUNK	DS<2 BLO	EYB	BKD	сор	DS>2	PRD	FID HBO	BO	BVT	HBP	TEA	ОРD	NIN	FVB	POP	GBD
Breitenbush River																		
5 ft	325	1	2						3				1	1	1			
Parr	10		2						2				1					
Fry	315	1							1					1	1			
Detroit HOR																		1
5 ft	559	2	5	5	1	1	4	2	10		1		52	4	5	2	2	1
Parr	10		2						3									
Smolt	2		1		1	1	1	1	1				1		1	1	1	
Fry	547	2	2	5			3	1	6		1		5 1	4	4	1	1	1
Green Peter HOR																		
5 ft	1																	
Fry	1																	


Steelhead (O. mykiss) During Reporting Period

	0.	mykiss Inj	uries Du	ring Repo	rting Peri	od 8-1-202	23 to 8-1	5-2023				
Site/Trap/Life Stage	Total Fish	DS<2 BLO	EYB FUN	BKD COP	DS>2 PRD	FID HBO	BO HO	BVT HBP	BRU TEA	OPD HIN	FVB POP	GBD
Breitenbush River												
5 ft	3											
Fry	1											
Parr	2											
Detroit HOR												
5 ft	12					1						
Fry	8					1						
Parr	4											

Appendix B Detroit Head of Reservoir-North Santiam River Operational and Capture Data Since Start of Monitoring







5-Aug

28-Jul

13-Aug

Breitenbush River Operational and Capture Data Since Start of Monitoring

0

1-May

9-May

17-May 25-May

2-Jun

10-Jun

18-Jun

26-Jun

Date


4-Jul


12-Jul


20-Jul



Green Peter Head of Reservoir-Middle Santiam River Operational and Capture Data Since Start of Monitoring







Hills Creek Head of Reservoir-Middle Fork Willamette River Operational and Capture Data Since Start of Monitoring

Appendix C

Release Location	Date of Release	# of Fish Released	# of Fish Recaptured	% Efficiency
Breitenbush River	6/21/2023	749	53	7.1%
Breitenbush River	7/6/2023	763	25	3.3%
Breitenbush River	8/2/2023	791	12	1.5%
Detroit Head of Reservoir- North Santiam River	6/6/2023	540	28	5.0%
Detroit Head of Reservoir- North Santiam River	6/20/2023	750	61	4.6%
Detroit Head of Reservoir- North Santiam River	7/6/2023	750	13	1.7%
Detroit Head of Reservoir- North Santiam River	8/2/2023	750	19	2.5%
Green Peter Head of Reservoir- Middle Santiam	6/7/2023	1,000 (dead fish)	0	0%
Green Peter Head of Reservoir- Middle Santiam	6/7/2023	750	1	0.1%
Green Peter Head of Reservoir- Middle Santiam	7/28/2023	750	0	0%
Lookout Dam Powerhouse*	4/13/2022	998	0	0%
Lookout Dam Powerhouse*	5/23/2023	3,999	32	0.8%
Lookout Dam Powerhouse*	6/1/2023	4,011	6	0.1%
Lookout Dam Powerhouse*	6/14/2023	4,010	4	0.1%
Lookout Dam Powerhouse*	6/28/2023	4,010	3	0.1%
Lookout Dam Powerhouse*	7/18/2023	4,012	1	0.02%
Hills Creek Head of Reservoir- Middle Fork Willamette	5/18/2023	519	44	8.5%
Hills Creek Head of Reservoir- Middle Fork Willamette	6/19/2023	760	7	0.9%

*Releases performed under the USACE RST contract

Appendix D

Summary of PIT Tagged Fish for Reporting Period

Site	Тгар	Species	# of PIT Tagged Fish
Breitenbush River	5 ft	Chinook	7
Breitenbush River	5 ft	O. mykiss	0
Detroit Head of Reservoir – North Santiam River	5 ft	Chinook	4
Detroit Head of Reservoir – North Santiam River	5 ft	O. mykiss	0
Green Peter Head of Reservoir – Middle Santiam River	5 ft	Chinook	0
Green Peter Head of Reservoir – Middle Santiam River	5 ft	O. mykiss	0
Lookout Dam Tailrace	Spill	Chinook	0
Lookout Dam Tailrace	PWR	Chinook	0
Hills Creek Head of Reservoir	5 ft	Chinook	0

Summary of EAS VIE Marked Fish for Reporting Period

Site	Trap	VIE Mark Code	Species	# VIE
Breitenbush River	5 ft	HYY	Chinook	1
Breitenbush River	5 ft	HYY	O. mykiss	3
Detroit Head of Reservoir – North Santiam River	5 ft	RDYY	Chinook	25
Detroit Head of Reservoir – North Santiam River	5 ft	RDYY	O. mykiss	7
Green Peter Head of Reservoir – Middle Santiam River	5 ft	RDYY	Chinook	0
Green Peter Head of Reservoir – Middle Santiam River	5 ft	RDYY	O. mykiss	0
Lookout Dam Tailrace	Spill	PYY	Chinook	0
Lookout Dam Tailrace	PWR	PYY	Chinook	0
Hills Creek Head of Reservoir	5 ft	LDYY	Chinook	0

RDYY denotes location and color (Right Dorsal Yellow (two stripes))

List of Captured Fish Containing PIT Tags This Season

Site	Trap	PIT Tag	Date	Species
N/A	N/A	N/A	N/A	N/A

List of EAS PIT Tagged Fish for Reporting Period

Site	Trap	PIT Tag	Date	Species
Breitenbush River	5 ft	3DD.003BEE0FB0	8/2/2023	Chinook
Breitenbush River	5 ft	3DD.003BEE1046	8/7/2023	Chinook
Breitenbush River	5 ft	3DD.003BEE103F	8/8/2023	Chinook
Breitenbush River	5 ft	3DD.003BEE1010	8/8/2023	Chinook
Breitenbush River	5 ft	3DD.003BEE102F	8/9/2023	Chinook

Breitenbush River	5 ft	3DD.003BEE1007	8/13/2023	Chinook
Breitenbush River	5 ft	3DD.003BEE0FBE	8/13/2023	Chinook
Detroit Head of Reservoir- North Santiam River	5 ft	3DD.003BEE1133	8/4/2023	Chinook
Detroit Head of Reservoir- North Santiam River	5 ft	3DD.003BEE1AE1	8/6/2023	Chinook
Detroit Head of Reservoir- North Santiam River	5 ft	3DD.003BEE0FE2	8/10/2023	Chinook
Detroit Head of Reservoir- North Santiam River	5 ft	3DD.0077CF7449	8/13/2023	Chinook