Dillon Alegre, Grant Brink & Rachel Ellison, Environmental Assessment

Prepared by: Services, LLC

Report Period: October 1st to October 15th, 2022

Report No.: 2022 Willamette RST Bi-Weekly Report 10/01-10/15 by EAS

Re: WILLAMETTE VALLEY FISH PASSAGE MONITORING VIA ROTARY

SCREW TRAPS

Project Schedule

Table 1. Project Schedule

| Site | Task | Start | End | Days |
|--|--|------------|------------|------|
| Big Cliff Dam RST | Operation | 12/01/2021 | 02/15/2022 | 000 |
| Big Cliff Dam RST | Operation | 03/15/2022 | 10/15/2022 | 292 |
| Big Cliff Dam Tailrace | Trap Efficiency Release (1,000 Fish) | 12/22/2021 | 12/22/2021 | 1 |
| Big Cliff Dam Tailrace | Temporary Trap Removal and Install | 05/06/2022 | 05/13/2022 | 7 |
| Big Cliff Dam Tailrace | Trap Efficiency Release (1,000 Fish) | 05/25/2022 | 05/25/2025 | 1 |
| Big Cliff Dam Tailrace | Trap Efficiency Release (1,000 Fish) | 08/09/2022 | 08/09/2022 | 1 |
| Big Cliff Dam Tailrace | Trap Efficiency Release (1,000 Fish) | 09/30/2022 | 09/30/2022 | 1 |
| Big Cliff Dam Tailrace | Trap Efficiency Release (500 Fish) | 10/13/2022 | 10/13/2022 | 1 |
| Green Peter Tailrace- Middle Santiam River RST | Trap Install | 03/02/2022 | 03/02/2022 | 1 |
| Green Peter Tailrace- Middle Santiam River RST | Operation | 03/03/2022 | 06/30/2022 | 120 |
| Green Peter Tailrace- Middle Santiam River RST | Trap Efficiency Release (643 Fish) | 03/29/2022 | 03/29/2022 | 1 |
| Green Peter Tailrace- Middle Santiam River RST | Trap Efficiency Release (521 Fish) | 04/30/2022 | 04/30/2022 | 1 |
| Green Peter Tailrace- Middle Santiam River RST | Temporary Trap Removal | 05/12/2022 | 05/12/2022 | 1 |
| Foster Dam Head of Reservoir- South Santiam River RST | Trap Install | 03/16/2022 | 03/16/2022 | 1 |
| Foster Dam Head of Reservoir- South Santiam River RST | Operation | 03/10/2022 | 06/30/2022 | 113 |
| Foster Dam Head of Reservoir- South Santiam River RST | Trap Removal | 07/01/2022 | 07/01/2022 | 1 |
| Foster Dam Head of Reservoir- South Santiam River RST | Trap Install | 09/02/2022 | 09/02/2022 | 1 |
| Foster Dam Head of Reservoir- South Santiam River RST | Trapping Efficiency Release (1000 fish) | 09/29/2022 | 09/29/2022 | 1 |
| Cougar Dam RST | Operation | 11/30/2021 | 11/30/2022 | 366 |
| Cougar Dam | Trap Efficiency Release (1,200 Fish, 600 per route) | 01/19/2022 | 01/19/2022 | 1 |
| Cougar Dam | Trap Efficiency Release (735 Fish, ~365 per route) | 04/20/2022 | 04/20/2022 | 1 |
| Cougar Dam | Trap Efficiency Release | 05/15/2022 | 05/15/2022 | 1 |

| | (993 Fish, RO route) | | | |
|--|---|------------|------------|-----|
| Cougar Dam | Trap Efficiency Release (500 Fish, PH route) | 07/19/2022 | 07/19/2022 | 1 |
| Cougar Dam | Trap Efficiency Release (501 Fish, PH route) | 08/11/2022 | 08/11/2022 | 1 |
| Cougar Dam | Trap Efficiency Release (442 Fish, RO route) | 10/14/2022 | 10/14/2022 | 1 |
| Cougar Dam Head of Reservoir | Highline and Trap Install | 03/07/2022 | 03/07/2022 | 1 |
| Cougar Dam Head of Reservoir | Operation | 03/08/2022 | 06/30/2022 | 115 |
| Cougar Dam Head of Reservoir | Trap Efficiency Release (806 Fish) | 03/18/2022 | 03/18/2022 | 1 |
| Cougar Dam Head of Reservoir | Trap Efficiency Release (500 Fish) | 05/19/2022 | 05/19/2022 | 1 |
| Cougar Dam Head of Reservoir | Trap Efficiency Release (515 Fish) | 06/23/2022 | 06/23/2022 | 1 |
| Cougar Dam Head of Reservoir | Trap Removal | 07/01/2022 | 07/01/2022 | 1 |
| Cougar Dam Head of Reservoir | Highline Install | 09/14/2022 | 09/14/2022 | 1 |
| Cougar Dam Head of Reservoir | Trap Install | 09/16/2022 | 09/16/2022 | 1 |
| Cougar Dam Head of Reservoir | Trap Efficiency Release (551 Fish) | 09/22/2022 | 09/22/2022 | 1 |
| Cougar Dam Head of Reservoir | Trap Efficiency Release (608 Fish) | 10/5/22 | 10/5/22 | 1 |
| Dexter Dam Tailrace RST | Highline Install | 03/02/2022 | 03/02/2022 | 1 |
| Dexter Dam Tailrace RST | Trap Install | 03/03/2022 | 03/03/2022 | 1 |
| Dexter Dam Tailrace RST | Operation | 03/07/2022 | 12/16/2022 | 285 |
| Dexter Dam Tailrace RST | Trap Efficiency Release (988 Fish) | 03/23/2022 | 03/23/2022 | 1 |
| Dexter Dam Tailrace RST | Trap Efficiency Release (1000 Fish) | 05/04/2022 | 05/04/2022 | 1 |
| Dexter Dam Tailrace RST | Trap Efficiency Release (1019 Fish) | 05/24/2022 | 05/24/2022 | 1 |
| Dexter Dam Tailrace RST | Trap Efficiency Release (981 Fish) | 07/21/2022 | 07/21/2022 | 1 |
| Lookout Dam Tailrace RSTs | Operation | 03/15/2022 | 07/31/2022 | 139 |
| Lookout Dam Tailrace RSTs | Trap Efficiency Release (1,013 fish, PWR route) | 04/13/2022 | 04/13/2022 | 1 |
| Lookout Point Head of Reservoir RST | Trap Install | 03/06/2022 | 03/06/2022 | 1 |
| Lookout Point Head of Reservoir RST | Operation | 03/07/2022 | 12/16/2022 | 285 |
| Lookout Point Head of Reservoir RST | Trap Efficiency Release (993 fish) | 04/05/2022 | 04/05/2022 | 1 |
| Lookout Point Head of Reservoir RST | Trap Efficiency Release (989 fish) | 04/14/2022 | 04/14/2022 | 1 |
| Lookout Point Head of Reservoir RST | Trap Efficiency Release (1007 fish) | 05/18/2022 | 05/18/2022 | 1 |
| Lookout Point Head of Reservoir RST | Trap Efficiency Release (1005 fish) | 07/20/2022 | 07/20/2022 | 1 |
| Fall Creek Dam Tailrace RST | Operation | 03/15/2022 | 07/15/2022 | 123 |
| Fall Creek Dam Tailrace RST | Trap Efficiency Release (518 fish) | 06/08/2022 | 06/08/2022 | 1 |
| Fall Creek Dam Tailrace RST | Trap Efficiency Release (513 fish) | 06/30/2022 | 06/30/2022 | 1 |
| Fall Creek Dam Tailrace RST | Trap Efficiency Release (500 fish) | 07/13/2022 | 07/13/2022 | 1 |
| Fall Creek Dam Tailrace RST | Deployment | 10/15/2022 | 10/15/2022 | 1 |

| Fall Creek Head of Reservoir RST | Trap and Highline Install | 01/11/2022 | 01/11/2022 | 1 |
|----------------------------------|--|------------|------------|-----|
| Fall Creek Head of Reservoir RST | Operation | 01/02/2022 | 05/31/2022 | 150 |
| Fall Creek Head of Reservoir RST | Removal | 06/02/2022 | 06/02/2022 | 1 |
| Hills Creek Dam RO and PWR | Deployment | 10/12/2021 | 10/12/2021 | 1 |
| Hills Creek Dam RO | Operation | 10/15/2021 | 03/01/2022 | 138 |
| Hills Creek Dam PWR | Operation | 10/15/2021 | 03/01/2022 | 138 |
| Hills Creek Dam | Trap Efficiency Release (1,200 fish, 600 per route) | 01/6/2022 | 01/6/2022 | 1 |
| Hills Creek Dam | Trap Efficiency Release (1,200 fish, 600 per route) | 02/16/2022 | 02/16/2022 | 1 |
| Hills Creek Dam | Trap Efficiency Release (1,200 fish, 600 per route) | 02/23/2022 | 02/23/2022 | 1 |
| Hills Creek Dam RSTs | Trap Removal | 03/01/2022 | 03/01/2022 | 1 |
| Hills Creek Dam RSTs | Trap Install | 09/14/2022 | 09/14/2022 | 1 |
| Hills Creek Dam RSTs | Deployment | 9/15/2022 | 9/15/2022 | 1 |

Summary of Rotary Screw Trap Data

Rotary screw traps (RSTs) have been operated at eleven locations in the southern Willamette River watershed. For this reporting period, traps were operated at the following eight locations: Big Cliff Dam, Foster Dam Head of Reservoir- South Santiam, Cougar Dam, Cougar Dam Head of Reservoir, Dexter Dam Tailrace, Lookout Dam Tailrace, Lookout Point Head of Reservoir, and Hills Creek Dam. The trap in the Fall Creek Dam RO channel resumed sampling on October 15th. The Green Peter Dam Tailrace-Middle Santiam trap was removed on May 12th due to damage incurred to the highline.

The RST's at Big Cliff Dam and Lookout Dam Tailrace started sampling on March 15th. On March 1st the Middle Fork Willamette River below Hills Creek Dam RST's were removed for the sampling season in conjunction with the end of RO spill and to prioritize the limited number of screw traps to other locations. Sampling at Hills Creek Dam resumed on September 15th.

Below Dam sites that include both RO and PWR to monitor passage routes include South Fork McKenzie River below Cougar Dam and on the Middle Fork of the Willamette River in the Lookout Dam Tailrace, and Hills Creek Dam. Below dam sites that include one RST to monitor passage include Big Cliff Dam, Green Peter Tailrace- Middle Santiam, the Middle Fork of the Willamette River below Dexter Dam and Fall Creek Dam Tailrace which is a tributary on the Middle Fork of the Willamette. At the Green Peter Dam Tailrace, the single RST is placed to sample fish passing through spillways, regulating outlets, and powerhouse outlets. The RST at Dexter Dam is placed to monitor fish passage through the spillways and powerhouse outlets. The RST in the Fall Creek Dam Tailrace is placed in a position to sample fish passing through the regulating outlet.

The RST on the North Santiam River below Big Cliff Dam was not sampled while fish passage measures were not being implemented from 16 February 2022 to 14 March 2022. Sampling resumed on 15 March 2022 in accordance with Task 2.2.

Above reservoir sites include Fall Creek Head of Reservoir, Lookout Point Head of Reservoir on the Middle Fork Willamette River, Foster Dam Head of Reservoir- South Santiam, and Cougar Dam Head of Reservoir on the South Fork McKenzie.

Sampling start dates are included in Table 2, and season total collection numbers are displayed in Table 3. The locations of the RST's are depicted in Figures 1 through 8.



Figure 1. Big Cliff RST Location



Figure 2. Foster Dam Head of Reservoir- South Santiam RST Location




Figure 3. Cougar Dam RST Locations



Figure 4. Cougar Dam Head of Reservoir RST Location




Figure 5. Fall Creek Dam Tailrace RST Location



Figure 6. Dexter Dam RST Location



Figure 7. Lookout Point Dam Tailrace RST Location



Figure 8. Lookout Point Head of Reservoir RST Location



Figure 9. Hills Creek Dam RST Locations

Table 2. Sampling Dates for Reporting Period

| Site | Total Sampling Period Start | Current Reporting Period Start | Current Reporting Period End | Days Sampled This Period | Total Days Sampled |
|---|-----------------------------------|-----------------------------------|---------------------------------|-----------------------------|-----------------------|
| Big Cliff Dam | 12/1/2021 | 10/01/2022 | 10/15/2022 | 15 | 274 |
| Foster Dam Head of Reservoir- South Santiam | 3/16/2022 | 10/01/2022 | 10/15/2022 | 15 | 144 |
| Cougar Dam PH | 12/1/2021 | 10/01/2022 | 10/15/2022 | 7 | 302 |
| Cougar Dam RO | 12/1/2021 | 10/01/2022 | 10/15/2022 | 15 | 312 |
| Cougar Dam Head of Reservoir | 3/7/2022 | 10/01/2022 | 10/15/2022 | 15 | 129 |
| Fall Creek Dam Tailrace* | 3/15/2022 | 10/01/2022 | 10/15/2022 | 0 | 123 |
| Dexter Dam Tailrace | 3/7/2022 | 10/01/2022 | 10/15/2022 | 10 | 214 |
| Lookout Point Dam PH | 3/15/2022 | 10/01/2022 | 10/15/2022 | 10 | 191 |
| Lookout Point Dam Spill | 3/15/2022 | 10/01/2022 | 10/15/2022 | 10 | 191 |
| Lookout Point Head of Reservoir | 3/10/2022 | 10/01/2022 | 10/15/2022 | 10 | 207 |
| Hills Creek Dam | 9/16/2022 | 10/01/2022 | 10/15/2022 | 10 | 23 |

^{*}Fall Creek Dam Tailrace trap was being operated by the Corps until EAS began sampling the site on March 15th per Task 7.1

Table 3. Willamette Valley Rotary Screw Trap Monitoring Catch Summary

| Site | Species | Catch (Reporting Period) | Recaptures (Reporting Period) | Total Catch | Total Recaptures |
|---|---------|--------------------------------|-------------------------------------|----------------|---------------------|
| Big Cliff Dam | CHS | 19 | 62 | 1106 | 215 |
| Big Cliff Dam | STW | 2 | 0 | 62 | 0 |
| Green Peter Tailrace- Middle Santiam | CHS | 0 | 0 | 0 | 13 |
| Green Peter Tailrace- Middle Santiam | STW | 0 | 0 | 6 | 0 |
| Foster Dam Head of Reservoir- South Santiam | CHS | 0 | 0 | 66 | 0 |
| Foster Dam Head of Reservoir- South Santiam | STW | 0 | 0 | 90 | 1 |
| Cougar Dam | CHS | 258 | 46 | 1619 | 438 |
| Cougar Dam Head of Reservoir | CHS | 13 | 47 | 632 | 174 |
| Fall Creek Dam Tailrace | CHS | 0 | 0 | 0 | 11 |
| Dexter Dam Tailrace | CHS | 0 | 0 | 97 | 113 |
| Lookout Point Dam | CHS | 0 | 0 | 78 | 2 |
| Lookout Point Head of Reservoir | CHS | 0 | 0 | 103 | 206 |
| Hills Creek Dam | CHS | 1 | 0 | 101 | 0 |

North Santiam – Big Cliff Dam

The RST on the North Santiam River below Big Cliff Dam was not sampled while fish passage measures were not being implemented from February 16th, 2022 to March 14th, 2022. Sampling resumed on March 15th, 2022 in accordance with Task 2.2.

Target Species

This reporting period began on October 1st and ended on October 15th. There were a total of 19 Chinook Salmon (CHS) and 2 Winter Steelhead (STW) captured during the 15-day sampling period (Figure 10). Sampling duration was 100% for the RST. Table 4 provides life stage, length, and weight data for all Chinook Salmon and Winter Steelhead that have been caught at the Big Cliff Dam site to-date and for the reporting period. Figure 11 shows length frequency data to-date.





Figure 10. Chinook and Winter Steelhead Captured per day 10/01/2022 to 10/15/2022 (Big Cliff)



*Figure does not include fish without heads

Figure 11. Length Frequency of Juvenile Chinook and Winter Steelhead Sampled Season To-Date (Big Cliff)

Trapping Efficiency

A total of 500 juvenile hatchery Chinook (sub-yearlings) were bismarck brown dyed, adipose clipped and released on 10/13/2022 below Big Cliff Dam. A total of 14 fish were recaptured in the 8ft trap. Trapping efficiency was 2.8%.

Trapping efficiency fish displayed minor descaling and fin damage.

| Big Cliff Dam | Release # | Recapture # | Capture Efficiency |
|---------------|-----------|-------------|-----------------------|
| 8ft Trap | 500 | 14 | 2.8% (14/500) |

Table 4. Descriptive Statistics of Target Species Captured at Big Cliff Dam Season To-Date

| | To-Date (Since Dec. 01, 2021) | | | | | | | | | | | |
|--------------|-------------------------------|---------|-------|-----------|-----|-----------|-------|------|----------|-------|--|--|
| Site | Route | Species | Life | Collected | L | ength (mr | m)* | | Weight (| g)* | | |
| Site | Route | Species | stage | Collected | Min | Max | Mean | Min | Max | Mean | | |
| | | CHS | Fry | 9 | 31 | 48 | 37.5 | 1.2 | 1.4 | 1.4 | | |
| | | CHS | Parr | 24 | 58 | 136 | 98.3 | 2.1 | 30.0 | 11.6 | | |
| Big Cliff | PWR | CHS | Smolt | 1071 | 91 | 283 | 142.3 | 8.9 | 253.5 | 33.8 | | |
| | | STW | Fry | 28 | 21 | 69 | 40.4 | 1.1 | 3.4 | 1.6 | | |
| | | STW | Parr | 17 | 51 | 111 | 65.9 | 1.3 | 14.3 | 3.9 | | |
| | | STW | Smolt | 17 | 157 | 284 | 214.1 | 36.1 | 230.5 | 102.2 | | |

*Fish that were missing heads are not included in length and weight calculations.

| | October 1-15, 2022 | | | | | | | | | | | |
|--------------|--------------------|---------|-------|-----------|-----|-----------|-------|------|------------|------|--|--|
| Site | Doute | Species | Life | Collected | | Length (m | nm)* | , | Weight (g) | * | | |
| Site | Route | Species | stage | Collected | Min | Max | Mean | Min | Max | Mean | | |
| | | CHS | Fry | 0 | N/A | N/A | N/A | N/A | N/A | N/A | | |
| | | CHS | Parr | 0 | N/A | N/A | N/A | N/A | N/A | N/A | | |
| | | CHS | Smolt | 19 | 143 | 170 | 157.5 | 29.0 | 55.1 | 43.4 | | |
| Big Cliff | PWR | STW | Fry | 1 | 48 | 48 | 48.0 | 1.5 | 1.5 | 1.5 | | |
| Oiiii | | STW | Parr | 1 | 58 | 58 | 58.0 | 2.0 | 2.0 | 2.0 | | |
| | | STW | Smolt | 0 | N/A | N/A | N/A | N/A | N/A | N/A | | |

^{*}Fish that were missing heads are not included in length and weight calculations.

24-Hour Post Collection Holding Trial

18 Spring Chinook and 2 Winter Steelhead were captured during the current reporting period and held for 24 hours. 3 Chinook (16.7%) and 0 Winter Steelhead (0.0%) died in holding.

Injuries and Copepod Infection

Partial descaling <20% was observed in 16 of the 19 Chinook captured (84.2%), 3 displayed descaling >20% (15.8%), 12 displayed body injury (63.2%), 0 had eye injury (0.0%), 19 had copepods present in the branchial cavity (100.0%) and 9 had copepods on fins (47.4%). 0 Chinook displayed gas bubble disease (0.0%). There was 1 mortality (5.3%).

Partial descaling <20% was observed in 0 of the 2 Winter Steelhead captured (0.0%) and 0 displayed descaling >20% (0.0%), 2 displayed body injury (100%), 0 had eye injury (0.0%), 0 had copepods present in the branchial cavity (0.0%) and 0 had copepods on fins (0.0%). There were 0 mortalities (0.0%). Injury data is summarized in table 5.

Table 5. Number of Descaled, Bodily/Eye Injured, Copepod Infected and Dead Chinook Salmon and Winter Steelhead for Sampling Period. (Big Cliff Dam).

| Site | Species | # Fish Collected | # DSC* <20% | # DSC* >20% | # with Body Injuries | # with Eye Injuries | # with COP* In B.C. | # with COP* on Fins | Mortalities |
|------------------|---------------------|---------------------|----------------|----------------|----------------------------|---------------------------|---------------------------|---------------------------|-------------|
| Big Cliff Dam | Chinook | 19 | 16 | 3 | 12 | 0 | 19 | 9 | 1 |
| Big Cliff Dam | Winter Steelhead | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

Collected DNA and Scale Samples

For the reporting period, scales and DNA were collected from 14 Spring Chinook and 2 Winter Steelhead. The other targets captured did not meet length criteria for DNA sampling or were captured before new protocols were put in place.

Non-Target Species

83 non-targets were captured during this sampling period. A summary of to-date non-target species catch and mortality numbers are listed in Table 6.

Table 6. Summary of Non-target Species (Big Cliff Dam)

| Species | PWR Capture | PWR Mortality | Season Total | Season Total Mortality |
|---------------------|----------------|------------------|-----------------|------------------------------|
| Bluegill | 59 | 0 | 123 | 10 |
| Bullhead | 0 | 0 | 5 | 0 |
| Chinook (Adult) | 0 | 0 | 2 | 1 |
| Chinook (clipped) | 0 | 0 | 9 | 0 |
| Cutthroat | 0 | 0 | 3 | 0 |
| Kokanee | 3 | 1 | 130 | 43 |
| O. mykiss (clipped) | 0 | 0 | 7 | 3 |
| Pumpkinseed | 21 | 0 | 97 | 5 |
| Unknown | 0 | 0 | 4 | 1 |
| Mountain Whitefish | 0 | 0 | 4 | 0 |
| Totals | 83 | 1 | 384 | 63 |

Stream Statistics

Basic stream statistics at the Big Cliff Dam site were calculated from data downloaded from U.S. Geological Survey stream gauge numbers 14181410 and 14181500. Gauge height (feet) is the only metric provided at gauge 14181410. Total dissolved gas (TDG) saturation data was received from gauge 14181500, 1 rkm downstream of the trap. During the reporting period, daily maximum values for instantaneous gauge height ranged from 1,109.6 to 1,111.8 feet (mean: 1,110.7 feet) during the reporting period. Figure 12 shows instantaneous gauge height.

Total dissolved gas saturation ranged from 99 to 122% during the reporting period (mean: 102.2%). Figure 113 shows total dissolved gas saturation.

Stream temperatures were recorded every 2 hours for the length of the report period at the RST (Figure 114). Temperature probes operated normally throughout this reporting period.

Flows through the Powerhouse and Spill during the reporting period averaged 2,295.5 and 44.3 cubic feet per second (cfs), respectively (Figure 15). Catch per unit of effort (CPUE) data are summarized in Table 7. Discharge and capture data for the duration of monitoring efforts at this location are provided in Appendix B.

Table 7. Summary of salmonid CPUE, Big Cliff Dam.

| | Chinook | Winter Steelhead |
|----------------|---------|------------------|
| Description | (8 ft) | (8 ft) |
| Catch | 19 | 2 |
| Effort (hrs) | 357.0 | 357.0 |
| CPUE (fish/hr) | 0.053 | 0.006 |



Figure 12. Gauge height (ft); below Big Cliff Dam




Figure 13. Total Dissolved Gas Saturation (%); below Big Cliff Dam




Figure 14. Temperature at RST (Big Cliff Dam)




Figure 15. Hourly Flows PWR vs. Spill (Big Cliff Dam)

South Fork Santiam- Foster Dam Head of Reservoir Target Species

This reporting period began on October 1st and ended on October 15th. There were 0 Chinook salmon (CHS) and 0 Winter Steelhead (STW) captured during the 15-day sampling period. Sampling duration was 100% for the RST. Table 8 provides life stage, length, and weight data for all Chinook Salmon and Winter Steelhead that have been caught at the Foster Dam Head of Reservoir- South Santiam site to-date and for the reporting period. Figure 16 shows the daily capture numbers for Chinook and Winter Steelhead and Figure 17 shows length frequency data to-date for both species.




Figure 16. Chinook and Winter Steelhead Captured Per Day 10/01/2022 to 10/15/2022 (Foster Dam Head of Reservoir- South Santiam)




Figure 17. Length Frequency of Juvenile Chinook and Winter Steelhead Sampled Season To-Date (Foster Dam Head of Reservoir- South Santiam)

Trapping Efficiency

A total of 1063 juvenile hatchery Chinook (sub-yearlings) were bismarck brown dyed, adipose clipped and released on 9/29/2022 at Cascadia Park above the Foster Dam Head of Reservoir- South Santiam trap. No fish have been recaptured to date.

| Foster Dam Head of Reservoir- South Santiam | Release | Recapture | Capture |
|---|---------|-----------|-------------|
| | # | # | Efficiency |
| 5 ft Trap | 1063 | 0 | 0% (0/1063) |

Run of River Trapping Efficiency

Run of river fish captured in the RST have been caudal clipped and released upstream to perform run of river trapping efficiency trials. Only fish large enough to be safely caudal clipped have been used for run of river efficiency trials. To date, 15 Chinook and 60 Winter Steelhead have been caudal clipped and released upstream for the purpose of conducting run of river trapping efficiency trials.

| Foster Dam Head of Reservoir- South Santiam | Release # | Recapture # |
|--|-----------|-------------|
| Chinook | 15 | 0 |
| Winter Steelhead | 60 | 1 |

| | To-Date (Since Dec. 01, 2021) | | | | | | | | | | | | |
|---------------------|-------------------------------|---------|---------------|-----------|--------------|-----|-------|-------------------------|------|------|--|--|--|
| Site | Trap | Species | Life stage | Collected | Length (mm)* | | | Weight (g) [*] | | | | | |
| Site | | | | Collected | Min | Max | Mean | Min | Max | Mean | | | |
| | 5 ft | CHS | Fry | 55 | 32 | 49 | 35.9 | N/A | N/A | N/A | | | |
| Foster | | CHS | Parr | 8 | 70 | 127 | 85 | 3.1 | 24.7 | 8.0 | | | |
| Dam Head of | | CHS | Smolt | 3 | 120 | 138 | 129 | 19.6 | 27.5 | 23.5 | | | |
| Reservoir- South | | STW | Fry | 31 | 28 | 46 | 34.6 | N/A | N/A | N/A | | | |
| Santiam | | STW | Parr | 17 | 65 | 183 | 116.7 | 2.4 | 63.6 | 20.9 | | | |
| | | STW | Smolt | 42 | 112 | 213 | 168.7 | 11.2 | 75.3 | 47.4 | | | |

| | October 1-15, 2022 | | | | | | | | | | | | |
|---|--------------------|---------|-------|-----------|--------------|-----|------|-----|-------------|------|--|--|--|
| Site | Trap | Species | Life | Collected | Length (mm)* | | | • | Weight (g)* | | | | |
| Site | | | stage | Collected | Min | Max | Mean | Min | Max | Mean | | | |
| | 5 ft | CHS | Fry | 0 | N/A | N/A | N/A | N/A | N/A | N/A | | | |
| _ , | | CHS | Parr | 0 | N/A | N/A | N/A | N/A | N/A | N/A | | | |
| Foster Dam | | CHS | Smolt | 0 | N/A | N/A | N/A | N/A | N/A | N/A | | | |
| Head of Reservoir- South Santiam | | STW | Fry | 0 | N/A | N/A | N/A | N/A | N/A | N/A | | | |
| | | STW | Parr | 0 | N/A | N/A | N/A | N/A | N/A | N/A | | | |
| | | STW | Smolt | 0 | N/A | N/A | N/A | N/A | N/A | N/A | | | |

Injuries and Copepod Infection

Partial descaling <20% was observed on 0 of the 0 Spring Chinook captured (0.0%). Body injuries were present on 0 Spring Chinook (0.0%) and 0 displayed eye injury (0.0%). No copepods were present on any of the Spring Chinook captured (0.0%). There were no mortalities.

Partial descaling <20% was observed on 0 of the 0 Winter Steelhead captured (0.0%). Body injuries were present on 0 Winter Steelhead (0.0%) and 0 displayed eye injury (0.0%). No copepods were present on any of the Winter Steelhead captured (0.0%). There were no mortalities. A summary of injuries observed during the reporting period are provided in Table 9, and for the duration of the season are provided in Appendix A.

Table 9. Number of Descaled, Bodily/Eye Injured, Copepod Infected and Dead Chinook Salmon and Winter Steelhead for Sampling Period. (Foster Dam Head of Reservoir- South Santiam).

| Site | Species | # Fish Collected | # DSC* <20% | # DSC* >20% | # with Body Injuries | # with Eye Injuries | # with COP* In B.C. | # with COP* on Fins | Mortalities |
|-------------------------------------|---------------------|---------------------|----------------|----------------|----------------------------|------------------------|---------------------------|---------------------------|-------------|
| Foster Dam Head of Reservoir- | Chinook | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| South Santiam | Winter Steelhead | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

Collected DNA and Scale Samples

For the reporting period, scales and DNA were collected from 0 Spring Chinook and 0 Winter Steelhead. The other targets captured did not meet length criteria for DNA sampling.

Non-Target Species

0 non-target species fish were captured during the reporting period; the data is summarized below in Table 10.

Table 10. Summary of Non-target Species (Foster Dam Head of Reservoir).

| Species | 5 ft Capture | 5 ft Mortality | Season Total | Season Total Mortality |
|-------------------|--------------|-------------------|-----------------|---------------------------|
| Dace | 0 | 0 | 42 | 1 |
| Sculpin | 0 | 0 | 2 | 1 |
| Largescale Sucker | 0 | 0 | 3 | 0 |
| Cutthroat | 0 | 0 | 28 | 0 |
| Unknown | 0 | 0 | 4 | 0 |
| Totals | 0 | 0 | 79 | 2 |

Stream Statistics

Basic stream statistics at the Foster Dam Head of Reservoir- South Santiam site were calculated from data downloaded from the U.S. Geological Survey stream gauge number 14185000. Discharge (cfs) and Gauge height (feet) are available at this gauge. During the reporting period, daily maximum values for instantaneous discharge ranged from 36.6 cfs to 58.7 cfs (mean: 42.3 cfs). Figure 18 shows instantaneous discharge.

Stream temperatures were recorded every 2 hours for the length of the report period for the RST (Figure 19). Temperature probes for the trap operated normally throughout this reporting period.

Catch per unit of effort (CPUE) data are summarized in Table 11. Discharge and capture data for the duration of monitoring efforts at this location are provided in Appendix B.

Table 11. Summary of salmonid CPUE, Foster Dam Head of Reservoir- South Santiam.

| | Chinook | Winter Steelhead | | | | |
|----------------|---------|------------------|--|--|--|--|
| Description | (5 ft) | | | | | |
| Catch | 0 | 0 | | | | |
| Effort (hrs) | 353.1 | 353.1 | | | | |
| CPUE (fish/hr) | 0 | 0 | | | | |




Figure 18. Discharge (cfs); Foster Dam Head of Reservoir – S. Santiam)





Figure 19. Temperature at RST (Foster Dam Head of Reservoir – S. Santiam)

South Fork McKenzie – Cougar Dam


Target Species

This reporting period began on October 1st and ended on October 15th. There were a total of 258 Chinook Salmon (CHS) captured during the 15-day sampling period. The flow in the powerhouse was lowered on 10/8/2022 in order to prioritize flow from the regulating outlet. The powerhouse traps were raised on 10/8/2022 in anticipation of this event to avoid damage to the cones. The flow is too low to continue fishing the powerhouse channel currently. Sampling duration was 100% for the RO RST and 46.7% for the Powerhouse RSTs. Table 12 provides life stage, length, and weight data for all Chinook salmon that have been caught at the Cougar Dam site to-date and for the reporting period. Figure 20 shows the daily capture numbers for chinook and Figure 21 shows length frequency data to-date.



*Recaptured fish for trapping efficiency trials not included.

Figure 20. Chinook Captured Per Day 10/01/2022 to 10/15/2022 (Cougar Dam)



*Figure does not include fish without heads or fish used for trapping efficiency trials.

Figure 21. Length Frequency of Juvenile Chinook Sampled Season To-Date (Cougar Dam)

Trapping Efficiency

A total of 442 juvenile hatchery Chinook (sub-yearlings) were bismarck brown dyed, adipose clipped, left ventral clipped and released on 10/14/2022. 37 fish were recaptured for an efficiency of 8.4%. Trapping

efficiency fish displayed injuries, primarily descaling, fin damage, and bloating. Hatchery staff noted that fish may have PKD at time of pickup.

| Cougar Dam | Release # | Recapture # | Capture Efficiency |
|------------|-----------|-------------|-----------------------|
| RO Route | 442 | 37 | 8.4% (37/442) |

Run of River Trapping Efficiency

Run of river fish were captured, caudal clipped and released for the purpose of conducting run of river trapping efficiency trials at Cougar Dam. Numbers of fish released and recaptured by route are reported below.

| Cougar Dam | Release # | Recapture # |
|------------|-----------|-------------|
| PH | 126 | 8 |
| RO | 20 | 1 |

Table 12. Descriptive Statistics of Target Species Captured at Cougar Dam Season To-Date

| | To-Date (Since Dec. 01, 2021) | | | | | | | | | | | | |
|------------------|-------------------------------|---------|-------|-----------|-----|-------------|-------|-------------|-------|------|--|--|--|
| Site | Route | Species | Life | Collected | ı | Length (mm) | * | Weight (g)* | | | | | |
| Site | | | stage | Collected | Min | Max | Mean | Min | Max | Mean | | | |
| Cougar Dam RO | | CHS | Fry | 21 | 34 | 48 | 40.8 | N/A | N/A | N/A | | | |
| | RO | CHS | Parr | 135 | 56 | 164 | 108.5 | 1.2 | 41.1 | 14.8 | | | |
| | | CHS | Smolt | 275 | 92 | 242 | 143.3 | 4.7 | 134.8 | 32.0 | | | |
| | | CHS | Fry | 382 | 25 | 55 | 38.0 | 1.0 | 1.8 | 1.2 | | | |
| Cougar Dam | PWR | CHS | Parr | 266 | 54 | 165 | 99.2 | 1.6 | 41.0 | 10.7 | | | |
| | | CHS | Smolt | 538 | 76 | 223 | 138.9 | 4.2 | 113.5 | 30.0 | | | |

*Fish that were missing heads are not included in length and weight calculations.

| | | | | Octo | ber 1-1 | 5, 2022 | | | | | |
|---------------|-------|---------|-------|-----------|---------|-------------|-------|------|-------------|------|--|
| 0.11 | 1 | Species | Life | 0.11() | ı | _ength (mm) | * | | Weight (g)* | | |
| Site Ro | Route | | stage | Collected | Min | Max | Mean | Min | Max | Mean | |
| | | CHS | Fry | 0 | N/A | N/A | N/A | N/A | N/A | N/A | |
| Cougar Dam | RO | CHS | Parr | 0 | N/A | N/A | N/A | N/A | N/A | N/A | |
| | | CHS | Smolt | 50 | 119 | 242 | 162.0 | 20.0 | 134.8 | 47.2 | |
| | | CHS | Fry | 0 | N/A | N/A | N/A | N/A | N/A | N/A | |
| Cougar Dam | PWR | CHS | Parr | 1 | 105 | 105 | 105.0 | 15.1 | 15.1 | 15.1 | |
| | | CHS | Smolt | 207 | 105 | 175 | 147.5 | 12.5 | 61.2 | 35.9 | |

*Fish that were missing heads are not included in length and weight calculations.

24-Hour Post Collection Holding Trial

A total of 111 Chinook captured in the RSTs, 71 fish from the PWR RST and 40 from the RO RST, were held for ~24 hours in holding tanks and then evaluated for survival rates. In total, 18 of the fish (16.2%) held during this period died during holding. 11 of the 71 PWR RST captured fish (15.5%) died during holding and 7 of the 40 RO RST captured fish (17.5%) died during holding.

Injuries and Copepod Infection

Partial descaling <20% was observed on 39 of the 50 Chinook collected at the RO RST (78.0%). Descaling >20% was observed on 10 of the Chinook (20.0%). There were 39 fish with bodily injuries (78.0%) and 4 had eye injuries (8.0%). 47 fish had copepods present in the branchial cavity (94.0%) and 37 had copepods present on fins (74.0%). 9 fish displayed Gas Bubble Disease (six level 1 and three Level 2) (18.0%). There were 5 chinook mortalities collected in the RO RST (10.0%).

Partial descaling <20% was observed on 141 of the 208 Chinook collected at the PWR RST (67.8%). Descaling >20% was observed on 36 of the Chinook (17.3%). There were 134 fish with bodily injuries (64.4%) and 7 had eye injuries (3.4%). 198 fish had copepods present in the branchial cavity (95.2%) and 143 had copepods present on fins (68.8%). 0 fish displayed Gas Bubble Disease (0.0%). There were 35

chinook mortalities collected in the PWR RST (16.8%). Data is summarized below in Table 13. A summary of injuries observed during the reporting period, and for the duration of the season are provided in Appendix A.

Table 13. Number of Descaled, Bodily/Eye Injured, Copepod Infected and Dead Chinook Salmon for Sampling Period. (Cougar Dam).

| Site | Route | # CHS Collected | # DSC* <20% | # DSC* >20% | # with Body Injuries | # with Eye Injuries | # with COP* In B.C. | # with COP* on Fins | Mortalities |
|--------|-------|--------------------|----------------|----------------|-------------------------|------------------------|---------------------------|---------------------------|-------------|
| Cougar | RO | 50 | 39 | 10 | 39 | 4 | 47 | 37 | 5 |
| Cougar | PWR | 208 | 141 | 36 | 134 | 7 | 198 | 143 | 35 |

^{*}DSC=Descaled, COP=Copepods, B.C.=Branchial Cavity

Collected DNA and Scale Samples

For the reporting period, scales and DNA were collected from 224 Spring Chinook. The other targets captured did not meet length criteria for DNA sampling or were captured before new protocols were put in place.

Non-Target Species

A total of 25 non-target species fish were captured during the reporting period; the data is summarized below in Table 14. Adipose clipped Chinook captured were from previous TE releases. They are not ODFW PIT tagged fish.

Table 14. Summary of Non-target Species (Cougar Dam).

| Species | RO Capture | RO Mortality | PWR Capture | PWR Mortality | Season Total Capture | Season Total Mortality |
|----------------------|---------------|-----------------|----------------|------------------|----------------------------|------------------------------|
| Brook Lamprey | 0 | 0 | 0 | 0 | 3 | 0 |
| Bluegill | 0 | 0 | 0 | 0 | 1 | 0 |
| Bull Trout | 0 | 0 | 0 | 0 | 1 | 0 |
| Chinook (AD clipped) | 3 | 0 | 3 | 1 | 9 | 1 |
| Chinook (Adult) | 0 | 0 | 0 | 0 | 1 | 0 |
| Cutthroat | 0 | 0 | 0 | 0 | 71 | 3 |
| Dace | 8 | 0 | 0 | 0 | 2502 | 8 |
| Largescale Sucker | 0 | 0 | 1 | 0 | 55 | 0 |
| Mountain Whitefish | 0 | 0 | 3 | 2 | 48 | 2 |
| Northern Pikeminnow | 0 | 0 | 0 | 0 | 2 | 0 |
| O. mykiss | 1 | 0 | 5 | 0 | 307 | 3 |
| Sculpin | 0 | 0 | 1 | 0 | 207 | 4 |
| Smallmouth Bass | 0 | 0 | 0 | 0 | 2 | 0 |
| Spotted Bass | 0 | 0 | 0 | 0 | 2 | 0 |
| Unknown | 0 | 0 | 0 | 0 | 21 | 2 |
| Totals | 12 | 0 | 13 | 3 | 3276 | 23 |

Stream Statistics

Basic stream statistics at the Cougar Dam site were calculated from data downloaded from U.S. Geological Survey stream gauge numbers 14159410 and 14181500. Gauge height (feet) is the only metric provided at gauge 14159410. Total dissolved gas saturation data was received from gauge 14181500, 500 meters downstream of the trap. During the reporting period, daily maximum values for instantaneous gauge height ranged from 1,251.9 to 1,253.9 feet (mean: 1,252.9 feet). Figure 22 shows instantaneous gauge height.

Total dissolved gas saturation ranged from 100 to 112% (mean: 105.9%). Figure 23 shows total dissolved gas saturation.

Stream temperatures were recorded every 2 hours for the length of the report period for the RO and PWR RST's (Figure 24 and 25 respectively). Temperature probes for the RO and PWR RST operated normally throughout this reporting period.

Flows through the Powerhouse and RO during the reporting period averaged 394.9 and 490.7 cubic feet per second (cfs) respectively (Figure 26). Catch per unit of effort (CPUE) data are summarized in Table 15. Discharge and capture data for the duration of monitoring efforts at this location are provided in Appendix B.

Table 15. Summary of salmonid CPUE, Cougar Dam.

| | Chinook | | | | |
|----------------|------------------|-------|--|--|--|
| Description | RO (5ft) PWR (8f | | | | |
| Catch | 50 | 208 | | | |
| Effort (hrs) | 362.5 | 338.5 | | | |
| CPUE (fish/hr) | 0.138 | 0.614 | | | |




Figure 22. Gauge Height (feet); below Cougar Dam, South Fork McKenzie River




Figure 23. Total Dissolved Gas Saturation (%); below Cougar Dam, South Fork McKenzie River




Figure 24. Temperature at RO RST (Cougar Dam)




Figure 25. Temperature at PWR RST (Cougar Dam)




Figure 26. Hourly Flows PWR vs. RO (Cougar Dam)

South Fork of the McKenzie-Cougar Dam Head of Reservoir

Target Species

The reporting period began October 1st and ended on October 15th. There were 13 Chinook salmon captured during the 15-day sampling period (Figure 27). The trap was operated 100% of the reporting period. Table 16 provides life stage, length, and weight data for all Chinook salmon that have been caught at the site to-date and Figure 28 shows length frequency data to-date.




Figure 27. Chinook Captured Per Day 10/01/2022 to 10/15/2022 (Cougar Dam Head of Reservoir)




Figure 28. Length Frequency of Juvenile Chinook Sampled Season To-Date (Cougar Dam Head of Reservoir)

Table 16. Descriptive Statistics of Target Species Captured at Cougar Dam Head of Reservoir, Season To-Date and for the Reporting Period

| To-Date (Since March 07, 2022) | | | | | | | | | | |
|--------------------------------|-------|---------|---------------|-----------|--------------------------|-----|------|-------------------------|------|------|
| Cito | Route | Species | Life stage | Collected | Length (mm) ⁻ | | | Weight (g) [.] | | |
| Site | Route | | | | Min | Max | Mean | Min | Max | Mean |
| Cougar | | CHS | Smolt | 4 | 70 | 94 | 82 | 3.3 | 9.4 | 6.2 |
| Dam Head of | 5 ft | CHS | Parr | 100 | 43 | 150 | 76.9 | 1.0 | 11.2 | 5.1 |
| Reservoir | | CHS | Fry | 528 | 27 | 63 | 38.6 | 0.6 | 2.8 | 1.4 |

| October 1-15, 2022 | | | | | | | | | | |
|------------------------------------|--------|-------------|---------------|-----------|--------------------------|-----|------|-------------------------|-----|------|
| Site | Davita | ute Species | Life stage | Collected | Length (mm) ⁻ | | | Weight (g) [,] | | |
| Site Rout | Route | | | | Min | Max | Mean | Min | Max | Mean |
| Cougar Dam Head of Reservoir | 5 ft | CHS | Smolt | 0 | N/A | N/A | N/A | N/A | N/A | N/A |
| | | CHS | Parr | 13 | 60 | 94 | 75.1 | 2.1 | 8.1 | 4.5 |
| | | CHS | Fry | 0 | N/A | N/A | N/A | N/A | N/A | N/A |

^{*}Most fry are too small to collect accurate weights and thus some metrics are not available for them.

Trapping Efficiency

A total of 608 juvenile hatchery Chinook (smolt) were adipose clipped, PIT tagged, and released on 10/5/2022 upstream of the Cougar Head of Reservoir trap site. A total of 47 fish were recaptured in the 5 ft trap. Trapping efficiency was 7.7%. Of the 47 fish recaptured, 38 displayed minor descaling and 41 had fin damage.

| Cougar Dam Head of Reservoir | Release # | Recapture # | Capture Efficiency |
|------------------------------|-----------|-------------|-----------------------|
| 5ft trap | 608 | 47 | 7.7% (47/608) |

Injuries and Copepod Infection

13 Chinook were captured for the reporting period. Of the fish captured, partial descaling <20% was observed on 4 fish (30.8%) and descaling >20% was observed on 1 fish (7.7%). 0 had copepods in the branchial cavity (0.0%), 1 had copepods on fins (7.7%), and 2 had bodily injury (15.4%). There were 2 mortalities for this reporting period (15.4%). These two fish did not recover from tagging. Injury data for the reporting period is summarized in Table 17. To date injury data can be found in Appendix A.

Table 17. Number of Descaled, Bodily/Eye Injured, Copepod Infected and Dead Chinook Salmon for Sampling Period. (Cougar Dam Head of Reservoir)

| Site | # CHS Collected | # DSC* <20% | # DSC* >20% | # with Body Injuries | # with Eye Injuries | # with COP* In B.C. | # with COP* on Fins | Mortalities |
|------------------------------------|--------------------|----------------|----------------|----------------------------|------------------------|------------------------|------------------------|-------------|
| Cougar Dam Head of Reservoir | 13 | 4 | 1 | 6 | 0 | 0 | 1 | 2 |

^{*}DSC=Descaled, COP=Copepods, B.C.=Branchial Cavity

Collected DNA and Scale Samples

Scales and DNA were collected from 13 of the Chinook captured. The rest of the captured fish were under the minimum fork length threshold and samples were not collected (less than 45 mm fork length for DNA and less than 50 mm fork length for scales).

VIE Marking

Visual Implant Elastomer (VIE) trials commenced at the Cougar Dam Head of Reservoir site on 6/25/2022. VIE tag color and locations are changed every reporting period to distinctly mark groups of fish by capture date. Since then, 32 Chinook have been VIE marked to the left of their dorsal fin with fluorescent elastomer. No fish with VIE marks have been detected at downstream RST sites to date.

| Date Tagged | VIE Color | # Tagged | # Recaptured to Date |
|----------------------|-----------|----------|----------------------|
| 6/25/2022-7/15/2022 | Yellow | 30 | 0 |
| 9/15/2022-9/30/2022 | Orange | 1 | 0 |
| 10/1/2022-10/15/2022 | Pink | 1 | 0 |

Non-Target Species

A total of 11 non-target fish were captured at the Cougar Dam Head of Reservoir RST during the reporting period; the data is summarized below in Table 18.

Table 18. Summary of Non-target Species (Cougar Dam Head of Reservoir)

| Species | 5ft Capture | 5ft Mortality | Season Total | Season Total Mortality |
|--------------------|-------------|------------------|-----------------|---------------------------|
| Bull Trout | 0 | 0 | 2 | 0 |
| Cutthroat Trout | 0 | 0 | 41 | 1 |
| Dace | 0 | 0 | 8 | 0 |
| Sculpin | 2 | 0 | 5 | 1 |
| O. mykiss | 0 | 0 | 332 | 3 |
| Mountain Whitefish | 9 | 0 | 11 | 0 |
| Unknown | 0 | 0 | 9 | 0 |
| Totals | 11 | 0 | 408 | 5 |

Stream Statistics

Basic stream statistics at the site were calculated from data downloaded from the U.S. Geological Survey stream gauge number 14159200. During the reporting period, daily maximum values for instantaneous discharge ranged from 200.0 cfs to 213.0 cfs (mean: 206.3 cfs). Figure 29 shows instantaneous discharge.

Stream temperatures were recorded every two hours using a temperature probe at the Cougar Dam Head of Reservoir RST site during this reporting period. Temperature probes operated normally, and the data is shown below in Figure 30.

Catch per unit of effort (CPUE) data are summarized in Table 19. Discharge and capture data for the duration of monitoring efforts at this location are provided in Appendix B.

Table 19. Summary of Chinook CPUE, Cougar Dam Head of Reservoir

| | Chinook |
|----------------|---------|
| Description | 5 ft |
| Catch | 13 |
| Effort (hrs) | 390.7 |
| CPUE (fish/hr) | 0.033 |




Figure 29. Discharge (cfs); South Fork McKenzie above Cougar Dam




Figure 30. Temperature at RST (Cougar Dam Head of Reservoir)


Fall Creek Dam Tailrace

The Fall Creek Dam Tailrace was installed and secured on October 14th in preparation to start fishing on October 15th per Task 7.7. The air quality in the Oakridge to Lowell corridor was in the Hazardous category and it was deemed unsafe for crews to operate. No Chinook have been captured at the Fall Creek Dam Tailrace trap to date.

Middle Fork Willamette- Dexter Dam


Target Species

This reporting period began on October 1st and ended on October 15th. There were 0 Chinook salmon (CHS) captured during the 15-day sampling period. The Dexter Dam trap was raised to a non-sampling position on October 10th due to hazardous air quality from the Cedar Creek Fire. The trap was not lowered to a sampling position again during this reporting period due to hazardous conditions. Sampling duration was 66.7% for the 5 ft RST. Table 20 provides life stage, length, and weight data for all Chinook salmon that have been caught at the Dexter Dam site to-date and for the reporting period. Figure 31 shows the daily capture numbers for Chinook and Figure 32 shows length frequency data to-date.



*Recaptured fish for trapping efficiency trials not included.

Figure 31. Chinook Captured Per Day 10/01/2022 to 10/15/2022 (Dexter Dam)



*Figure does not include fish without heads or fish used for trapping efficiency trials.

Figure 32. Length Frequency of Juvenile Chinook Sampled Season To-Date (Dexter Dam)

Table 20. Descriptive Statistics of Target Species Captured at the Dexter Dam RST Season To-Date

| | To-Date (Since March 07, 2022) | | | | | | | | | |
|---------------|--------------------------------|---------|-------|-----------|-----|----------|-------|-------------|-------|------|
| Site | Tron | Species | Life | e | Le | ngth (mm |)* | Weight (g)* | | |
| Site | Trap | Species | stage | Collected | Min | Max | Mean | Min | Max | Mean |
| | | CHS | Fry | 3 | 46 | 55 | 51.3 | 1.4 | 1.4 | 1.4 |
| Dexter Dam | 5 ft | CHS | Parr | 18 | 51 | 159 | 101.3 | 2.1 | 48.3 | 12.8 |
| | | CHS | Smolt | 77 | 95 | 224 | 137.2 | 9.3 | 118.4 | 27.6 |

| | October 1-15, 2022 | | | | | | | | | |
|---------------|--------------------|---------|-------|-----------|-----|-----------|------|-------------|-----|------|
| Site | - | 0 | Life | | Le | ength (mm |)* | Weight (g)* | | |
| | Trap | Species | stage | Collected | Min | Max | Mean | Min | Max | Mean |
| | | CHS | Fry | 0 | N/A | N/A | N/A | N/A | N/A | N/A |
| Dexter Dam | 5 ft | CHS | Parr | 0 | N/A | N/A | N/A | N/A | N/A | N/A |
| Daili | | CHS | Smolt | 0 | N/A | N/A | N/A | N/A | N/A | N/A |

^{*}Fish that were missing heads are not included in length and weight calculations.

Trapping Efficiency

A total of 981 juvenile hatchery Chinook (sub-yearlings) were bismarck brown dyed, adipose clipped, upper caudal clipped and released on 07/21/2022 below Dexter Dam. Fish were released in small groups into powerhouse flow to evaluate the traps efficiency capturing fish passing through the powerhouse. 1 fish was recaptured in the 5-foot RST for an efficiency of 0.1%.

Mt. Hood Environmental staff noted that fish appeared to be in good condition upon retrieval from the hatchery.

| Dexter Dam | Release # | Recapture # | Capture Efficiency |
|------------|-----------|-------------|-----------------------|
| Spill | N/A | N/A | N/A |
| Powerhouse | 981 | 1 | 0.1% (1/981) |

24-Hour Post Collection Holding Trial

No Chinook were captured this reporting period.

Injuries and Copepod Infection

0 Chinook was captured during this reporting period. Partial descaling <20% was observed in 0 of the 0 Chinook captured (0.0%) and 0 displayed descaling >20% (0.0%). 0 displayed body injury (0.0%) and 0 Chinook had eye injury (0.0%). 0 Chinook had copepods present in the branchial cavity (0.0%) and 0 had

copepods on fins (0.0%). 0 displayed gas bubble disease (0.0%). There were 0 mortalities this reporting period (0.0%). Injuries are displayed in Table 21. To date injury data can be found in Appendix A.

Table 21. Number of Descaled, Bodily/Eye Injured. Copepod Infected and dead Chinook Salmon for Sampling Period (Dexter Dam).

| Site | Route | # CHS Collected | # DSC* <20% | # DSC* >20% | # with Body Injuries | # with Eye Injuries | # with COP* In B.C. | # with COP* on Fins | Mortalities |
|---------------|-------|--------------------|----------------|----------------|-------------------------|------------------------|------------------------|------------------------|-------------|
| Dexter Dam | PWR | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

^{*}DSC=Descaled, COP=Copepods, B.C.=Branchial Cavity

Collected DNA and Scale Samples

For the reporting period, scales and DNA were collected from 0 Spring Chinook. The other targets captured did not meet length criteria for DNA sampling.

Non-Target Species

45 non-target species fish were captured during the reporting period; the data is summarized below in Table 22. Dexter Fish Facility releases adipose clipped Chinook near our trapping site. Adipose clipped Chinook caught in our trap are being counted as non-target fish to provide better clarity on data concerning run of river Chinook.

Table 22. Summary of Non-target Species (Dexter Dam).

| Species | Capture | Mortality | Season Total | Season Total Mortality |
|---------------------|---------|-----------|-----------------|------------------------------|
| Bass | 2 | 0 | 99 | 1 |
| Bluegill | 1 | 0 | 8 | 1 |
| Chinook (clipped) | 0 | 0 | 357 | 7 |
| Chinook (adult) | 0 | 0 | 2 | 2 |
| Crappie | 6 | 0 | 52 | 5 |
| Cutthroat | 0 | 0 | 3 | 0 |
| Dace | 0 | 0 | 31 | 6 |
| O. mykiss | 0 | 0 | 16 | 0 |
| O. mykiss (clipped) | 0 | 0 | 34 | 2 |
| Pikeminnow | 0 | 0 | 1 | 0 |
| Red-Sided Shiner | 0 | 0 | 3 | 0 |
| Sculpin | 36 | 1 | 447 | 15 |
| Sucker | 0 | 0 | 3 | 1 |
| Unknown | 0 | 0 | 10 | 4 |
| Totals | 45 | 1 | 1066 | 38 |

Stream Statistics

Basic stream statistics at the Dexter Dam site were calculated from data downloaded from the U.S. Geological Survey stream gauge numbers 14149510 and 14150000. Gauge height (feet) is the only

metric provided at gauge 14149510. Total dissolved gas saturation data was received from gauge 14150000, 4.75 rkms downstream of the trap. During the reporting period, daily maximum values for instantaneous gauge height ranged from 637.6 feet to 637.7 feet (mean: 637.6 feet). Figure 33 shows instantaneous gauge height.

Total dissolved gas saturation ranged from 98 to 110% (mean: 102.7%) during the reporting period. Figure 34 shows total dissolved gas saturation.

Stream temperatures were recorded every two hours using a temperature probe at the Dexter Dam RST site during this reporting period. Temperature probes operated normally, and the data is shown below in Figure 35.

Flows through the Powerhouse and Spill during the reporting period averaged 1,581.3 and 0 cubic feet per second (cfs) respectively (Figure 36). Catch per unit of effort (CPUE) data are summarized in Table 23. Discharge and capture data for the duration of monitoring efforts at this location are provided in Appendix B.

Table 23. Summary of salmonid CPUE, Dexter Dam.

| | <u> </u> |
|----------------|----------|
| | Chinook |
| Description | 8 ft |
| Catch | 0 |
| Effort (hrs) | 240.6 |
| CPUE (fish/hr) | 0 |




Figure 33. Gauge Height (feet); below Dexter Dam, Middle Fork Willamette




Figure 34. Total Dissolved Gas Saturation (%); Middle Fork Willamette River, Near Dexter, OR




Figure 35. Temperature at RST (Dexter Dam)




Figure 36. Hourly Flows PWR vs. Spill (Dexter Dam)

Middle Fork Willamette - Lookout Dam Tailrace

Target Species

The reporting period began October 1st and ended on October 15th. 0 Chinook salmon were captured during the 15-day sampling period (Figure 37). The Lookout Dam Tailrace traps were raised to a non-sampling position on October 10th due to hazardous air quality from the Cedar Creek Fire. They were not

lowered to a sampling position again during this reporting period due to hazardous conditions. The traps were operated 66.7% of the reporting period. Table 24 provides life stage, length, and weight data for all Chinook salmon that have been caught at the Lookout Point Dam Tailrace site to-date and Figure 38 shows length frequency data to-date.




Figure 37. Chinook Captured Per Day 10/01/2022 to 10/15/2022 (Lookout Point Dam Tailrace)




Figure 38. Length Frequency of Juvenile Chinook Sampled Season To-Date (Lookout Point Dam Tailrace)

Table 24. Descriptive Statistics of Target Species Captured at Lookout Point Dam Tailrace, Season To-Date and for the Reporting Period

| | | | To-Da | te (Since Ma | rch 15, : | 2022) | - | | | |
|----------------------|----------------|---------|-------|--------------|-----------|---------|---|------|----------|-----------------|
| Site | \Route | Species | Life | Callested | Le | ngth (m | nm)* | , | Weight (| g) [*] |
| Site | Oite Woute Ope | | stage | Collected | Min | Max | Mean | Min | Max | Mean |
| | | CHS | Smolt | 25 | 112 | 266 | 150.3 | 15.0 | 108.9 | 40.6 |
| | PH 1 | CHS | Parr | 3 | 84 | 107 | 94.3 | 3.8 | 10.5 | 6.5 |
| | | CHS | Fry | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | | CHS | Smolt | 8 | 95 | 141 | 119.6 | 8.4 | 32.3 | 19.9 |
| Lookout Point Dam | PH 2 | CHS | Parr | 4 | 58 | 108 | 86.0 | 2.2 | 13.4 | 6.7 |
| | | CHS | Fry | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | Spill | CHS | Smolt | 32 | 94 | 194 | 133.8 | 7.6 | 63.0 | 27.2 |
| | | CHS | Parr | 6 | 77 | 126 | 96.5 | 5.4 | 26.1 | 12.1 |
| | | CHS | Fry | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| | | | | October 1-15 | 5, 2022 | | | | | |
| Site | Route | Species | Life | Collected | Le | ngth (n | (mm) [*] Weight (g) [*] | | | g)* |
| Onc | Route | Opecies | stage | Collected | Min | Max | Mean | Min | Max | Mean |
| | | CHS | Smolt | 0 | N/A | N/A | N/A | N/A | N/A | N/A |
| | PH 1 | CHS | Parr | 0 | N/A | N/A | N/A | N/A | N/A | N/A |
| | | CHS | Fry | 0 | N/A | N/A | N/A | N/A | N/A | N/A |
| | | CHS | Smolt | 0 | N/A | N/A | N/A | N/A | N/A | N/A |
| Lookout Point Dam | PH 2 | CHS | Parr | 0 | N/A | N/A | N/A | N/A | N/A | N/A |
| i oint Dain | | CHS | Fry | 0 | N/A | N/A | N/A | N/A | N/A | N/A |
| | | CHS | Smolt | 0 | N/A | N/A | N/A | N/A | N/A | N/A |
| | \Spill | CHS | Parr | 0 | N/A | N/A | N/A | N/A | N/A | N/A |
| | (Op | CHS | Fry | 0 | N/A | N/A | N/A | N/A | N/A | N/A |

^{*}Some fry are too small to accurately weigh and are omitted from the above tables.

24-Hour Post Collection Holding Trial

0 Chinook captured in the RSTs was held during this reporting period. 0 fish were held from the PWR RST and 0 fish were held from the Spill RST. 0 hold fish died from the PWR RSTs (0 from PH 1 and 0 from PH 2) (0.0%). 0 of the fish from Spill RST died during holding (0.0%).

Trapping Efficiency

A total of 1,013 juvenile hatchery Chinook (parr) were bismarck brown dyed and adipose clipped, right ventral fin clipped and released on 04/13/2022 below Lookout Point Dam. Fish were released in small groups directly into powerhouse flow at 17:00 to 19:00. 2 fish were recaptured in the PH 1 RST for an

efficiency of 0.2%. 1 Lookout Point Dam trap efficiency fish was captured downstream in the Dexter RST on 4/15/2022.

Mt. Hood Environmental staff noted that fish appeared to be in good condition upon retrieval from the hatchery but did note some descaling and fin damage present as is common among hatchery fish of this age.

| Lookout Dam | Release # | Recapture # | Capture Efficiency |
|-------------|-----------|-------------|-----------------------|
| Powerhouse | 1,013 | 2 | 0.2% (2/1,013) |

Injuries and Copepod Infection

There were 0 Chinook captured in the Spill Channel RST. Partial descaling <20% was observed on 0 of 0 Chinook collected at the Spill RST (0.0%), and descaling >20% was observed on 0 of the Chinook collected (0.0%). 0 displayed body injuries (0.0%) and 0 had eye injuries (0.0%). None of the Spill RST Chinook had copepods present in the branchial cavity (0.0%) and 0 had copepods present on fins (0.0%). 0 of the fish captured in the Spill RST displayed Gas Bubble Disease (0.0%).

There were 0 Chinook captured in the Powerhouse channel RSTs. Partial descaling <20% was observed on 0 of the 0 Chinook collected at the PWR RSTs (0.0%). Descaling >20% was observed on 0 of the Chinook collected (0.0%). 0 PWR RST fish had bodily injury (0.0%) and 0 had eye injuries (0.0%). 0 of the fish had copepods present in the branchial cavity (0.0%) and 0 had copepods present on fins (0.0%). 0 fish displayed Gas Bubble Disease (0.0%). There were 0 chinook mortality collected in the Spill RST (0.0%) and 0 in the PWR RST (0.0%). Injuries are displayed in Table 25. To date injury data can be found in Appendix A.

Table 25. Number of Descaled, Bodily/Eye Injured, Copepod Infected and Dead Chinook Salmon for Sampling Period (Lookout Point Dam Tailrace).

| Site | Route | # CHS Collected | # DSC* <20% | # DSC* >20% | # with Body Injuries | # with Eye Injuries | # with COP* In B.C. | # with COP* on Fins | Mortalities |
|-------------------|-------|--------------------|----------------|----------------|----------------------------|---------------------------|---------------------------|---------------------------|-------------|
| Lookout Point Dam | PWR | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Tailrace | Spill | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

^{*}DSC=Descaled, COP=Copepods, B.C.=Branchial Cavity

Collected DNA and Scale Samples

For the reporting period, scales and DNA were collected from 0 Spring Chinook. The other targets captured did not meet length criteria for DNA sampling.

Non-Target Species

64 non-target species were captured during the reporting period; the data is summarized below in Table 26.

Table 26. Summary of Non-target Species (Lookout Point Dam Tailrace).

| | PWR | PWR | Spill | Spill | Season | Season Total |
|---------------------|---------|-----------|---------|-----------|--------|-----------------|
| Species | Capture | Mortality | Capture | Mortality | Total | Mortality |
| Bass Unknown | 3 | 2 | 0 | 0 | 4963 | 3662 |
| Bluegill | 0 | 0 | 0 | 0 | 3 | 0 |
| Bullhead | 0 | 0 | 0 | 0 | 3 | 1 |
| Chinook (clipped) | 0 | 0 | 0 | 0 | 4 | 0 |
| Crappie | 44 | 17 | 1 | 0 | 416 | 194 |
| Cutthroat | 0 | 0 | 0 | 0 | 1 | 0 |
| Largemouth Bass | 0 | 0 | 0 | 0 | 1 | 0 |
| Smallmouth Bass | 0 | 0 | 0 | 0 | 1 | 1 |
| Largescale Sucker | 0 | 0 | 0 | 0 | 25 | 15 |
| Northern Pikeminnow | 0 | 0 | 0 | 0 | 16 | 9 |
| O. mykiss | 0 | 0 | 0 | 0 | 8 | 1 |
| O. mykiss (clipped) | 0 | 0 | 0 | 0 | 2 | 1 |
| Red-Sided Shiner | 0 | 0 | 0 | 0 | 1 | 0 |
| Sculpin | 8 | 0 | 7 | 1 | 100 | 12 |
| Walleye | 0 | 0 | 0 | 0 | 10 | 3 |
| Unknown | 1 | 1 | 0 | 0 | 3 | 2 |
| Totals | 56 | 20 | 8 | 1 | 5474 | 3901 |

Stream Statistics

Basic stream statistics at Lookout Dam Tailrace site were calculated from data downloaded from the U.S. Geological Survey stream gauge number 14149010. Total dissolved gas saturation or dissolved oxygen concentration measurements are not available at this stream gauge site, or any nearby stream gauges. Gauge height (feet) is the only metric provided at this gauge. During the reporting period, daily maximum values for instantaneous gauge height ranged from 691.8 feet to 693.1 feet (mean: 692.4 feet). Figure 39 shows instantaneous gauge height.

Stream temperatures were recorded every two hours using a temperature probe at the Lookout Dam RST site during this reporting period. Temperature probes operated normally, and the data is shown below in (figures 40 and 41).

Flows through the Powerhouse and Spill during the reporting period averaged 1,118.0 and 566.7 cubic feet per second (cfs) respectively (Figure 42). Catch per unit of effort (CPUE) data are summarized in Table 27. Discharge and capture data for the duration of monitoring efforts at this location are provided in Appendix B.

Table 27. Summary of Chinook CPUE at Lookout Point Dam Tailrace.

| | Chinook | | | | | | |
|----------------|---------|-------|-------|--|--|--|--|
| Description | PH 1 | PH 2 | Spill | | | | |
| Catch | 0 | 0 | 0 | | | | |
| Effort (hrs) | 240.1 | 239.7 | 239.1 | | | | |
| CPUE (fish/hr) | 0 | 0 | 0 | | | | |




Figure 39. Gauge Height (feet); below Lookout Dam




Figure 40. Temperature at RST (Lookout Dam PWR)




Figure 41. Temperature at RST (Lookout Dam Spill)




Figure 42. Hourly Flows PWR vs. Spill (Lookout Dam Tailrace)

Middle Fork Willamette – Lookout Point Head of Reservoir Target Species

The reporting period began October 1st and ended on October 15th. 0 Chinook salmon were captured during the 15-day sampling period (Figure 43). The Lookout Point Head of Reservoir trap was raised on October 10th due to hazardous air quality conditions from the Cedar Creek fire. It was not returned to a sampling position during this reporting period due to hazardous conditions. The trap was operated 66.7% of the reporting period. Table 28 provides life stage, length, and weight data for all Chinook salmon that have been caught at the Middle Fork Willamette - Lookout Point Head of Reservoir site to-date and Figure 44 shows length frequency data to-date.




Figure 43. Chinook Captured Per Day 10/01/2022 to 10/15/2022 (Lookout Point Head of Reservoir)




Figure 44. Length Frequency of Juvenile Chinook Sampled Season To-Date (Lookout Point Head of Reservoir)

Table 28. Descriptive Statistics of Target Species Captured at Lookout Point Head of Reservoir, Season To-Date and for the Reporting Period

| | To-Date | | | | | | | | | | | |
|-----------------------------|---------|----------|-------|--------------|--------------|---------|-------|-------------|-------------|------|--|--|
| Site | Route | Species | Life | Collected | Le | ngth (m | ım)* | Weight (g)* | | | | |
| Site | Route | Species | stage | Collected | Min | Max | Mean | Min | Max | Mean | | |
| Lookout Point Head of | | CHS | Smolt | 3 | 111 | 118 | 114.5 | 15.0 | 17.5 | 16.3 | | |
| | 5 ft | CHS | Parr | 59 | 59 | 115 | 80.0 | 1.0 | 19.8 | 5.9 | | |
| Reservoir | | CHS | Fry | 41 | 28 | 69 | 37.2 | N/A | N/A | N/A | | |
| | | | | October 1-15 | , 2022 | | | | | | | |
| C:ta | Davita | Consider | Life | Callagead | Length (mm)* | | | | Weight (g)* | | | |
| Site | Route | Species | stage | Collected | Min | Max | Mean | Min | Max | Mean | | |
| Lookout | | CHS | Smolt | 0 | N/A | N/A | N/A | N/A | N/A | N/A | | |
| Point Head of | 5 ft | CHS | Parr | 0 | N/A | N/A | N/A | N/A | N/A | N/A | | |
| Reservoir | | CHS | Fry | 0 | N/A | N/A | N/A | N/A | N/A | N/A | | |

^{*}Some fry are too small to accurately weigh and are omitted from the above tables.

Trapping Efficiency

A total of 1,005 juvenile hatchery Chinook (sub-yearlings) were bismarck brown dyed and adipose clipped and released on 7/20/2022 above the Lookout Point Head of Reservoir trap. Fish were released in small groups to evaluate the traps efficiency capturing fish migrating downstream. 9 fish were recaptured in the 5-foot RST for an efficiency of 0.9%.

Of the 9 fish recaptured, 1 was dead. Injuries were descaling and fin damage.

| Lookout Point Head of Reservoir | Release # | Recapture # | Capture Efficiency | |
|------------------------------------|-----------|-------------|-----------------------|--|
| 07/20/2022 | 1005 | 9 | 0.9% (9/1005) | |

Injuries and Copepod Infection

There were 0 Chinook captured during this reporting period. 0 had partial descaling <20% (0.0%) and 0 had body injuries (0.0%). There was 0 incidental mortality (0.0%). Injury data for the reporting period is shown in table 29. To date data can be found in Appendix A.

Table 29. Number of Descaled, Bodily/Eye Injured, Copepod Infected and Dead Chinook Salmon for Sampling Period (Lookout Point Head of Reservoir).

| Site | # CHS Collected | # DSC* <20% | # DSC* >20% | # with Body Injuries | # with Eye Injuries | # with COP* In B.C. | # with COP* on Fins | Mortalities |
|--|--------------------|----------------|----------------|----------------------------|---------------------------|------------------------|------------------------|-------------|
| Lookout Point Head of Reservoir | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

^{*}DSC=Descaled, COP=Copepods, B.C.=Branchial Cavity

Collected DNA and Scale Samples

No scales and DNA were collected from Chinook captured for the reporting period.

VIE Marking

Visual Implant Elastomer (VIE) trials commenced at the Lookout Point Head of Reservoir site on 6/25/2022. VIE tag color and locations are changed every reporting period to distinctly mark groups of fish by capture date. Since then, 4 Chinook have been VIE marked to the left of the dorsal fin with fluorescent elastomer. No fish with VIE marks have been detected at downstream RST sites to date.

| Date Tagged | VIE Color | # Tagged | # Recaptured to Date | |
|---------------------|-----------|----------|----------------------|--|
| 6/25/2022-7/15/2022 | Yellow | 3 | 0 | |
| 7/16/2022-7/31/2022 | Red | 1 | 0 | |

Non-Target Species

6 non-target species were captured during the reporting period; the data is summarized below in Table 30.

Table 30. Summary of Non-target Species (Lookout Point Head of Reservoir).

| Species | 5ft Capture | 5ft Mortality | Season Total | Season Total Mortality |
|---------------------|----------------|------------------|-----------------|------------------------------|
| Bass Unknown | 0 | 0 | 2 | 0 |
| Bluegill | 0 | 0 | 2 | 0 |
| Chinook (clipped) | 0 | 0 | 26 | 0 |
| Cutthroat Trout | 0 | 0 | 10 | 0 |
| Dace | 2 | 0 | 127 | 0 |
| Lamprey | 0 | 0 | 2 | 0 |
| Largescale Sucker | 1 | 0 | 16 | 1 |
| Mountain Whitefish | 0 | 0 | 3 | 0 |
| Northern Pikeminnow | 0 | 0 | 24 | 0 |
| O. mykiss | 1 | 0 | 86 | 3 |
| O. mykiss (clipped) | 0 | 0 | 2 | 0 |
| Peamouth | 0 | 0 | 1 | 0 |
| Pumpkinseed | 1 | 1 | 1 | 1 |
| Red-Sided Shiner | 0 | 0 | 2 | 0 |
| Sculpin | 1 | 0 | 24 | 7 |
| Smallmouth Bass | 0 | 0 | 8 | 0 |
| Unknown | 0 | 0 | 10 | 0 |
| Totals | 6 | 1 | 338 | 12 |

Stream Statistics

Basic stream statistics for the Lookout Point Head of Reservoir RST site were calculated from data downloaded from the U.S. Geological Survey stream gauge number 14148000. During the reporting period, daily maximum values for instantaneous discharge ranged from 961.0 cfs to 1,170.0 cfs (mean: 1,129.4 cfs). Figure 45 shows instantaneous discharge.

Stream temperatures were recorded every two hours using a temperature probe at the Lookout Point Head of Reservoir RST site during this reporting period. The probe operated normally during this period. (Figure 46).

Flows into Lookout Point Reservoir averaged 1,153.4 cfs (Figure 47). Catch per unit of effort (CPUE) data are summarized in Table 31. Discharge and capture data for the duration of monitoring efforts at this location are provided in Appendix B.

Table 31. Summary of Chinook CPUE at Lookout Point Head of Reservoir.

| | Chinook |
|----------------|---------|
| Description | 5 ft |
| Catch | 0 |
| Effort (hrs) | 241.4 |
| CPUE (fish/hr) | 0 |




Figure 45. Discharge (cfs); above Lookout Point Reservoir, Below Oakridge, OR




Figure 46. Temperature at RST (Lookout Point Head of Reservoir)




Figure 47. Hourly Flows (Lookout Point Head of Reservoir site)

Middle Fork Willamette - Hills Creek Dam

Target Species

This reporting period began on October 1st and ended on October 15th. There was 1 Chinook salmon (CHS) captured during the 15-day sampling period (Figure 48). Cones were raised on October 10th due to hazardous air quality conditions from the Cedar Creek fire. The traps were not returned to sampling position during this reporting period due to hazardous conditions. Sampling durations were 66.7% for both the RO RST and Powerhouse RST. Table 4 provides life stage, length, and weight data for all Chinook salmon that have been caught at the Hills Creek Dam site to-date and Figure 49 shows length frequency data to-date.





Figure 48. Chinook Captured Per Day 10/01/2022 to 10/15/2022 (Hills Creek Dam)



*Figure does not include fish without heads or fish used for trapping efficiency

Figure 49. Length Frequency of Juvenile Chinook Sampled Season To-Date (Hills Creek Dam)

24-Hour Post Collection Holding Trial

0 Chinook captured in the RSTs were held during this reporting period. 0 fish were held from the PWR RST and 0 fish were held from the RO RST. 0 hold fish died from the PWR RST (0.0%). 0 of the fish from RO RST died during holding (0.0%).

1 Chinook was captured this reporting period but was not held for 24 hours as the air quality index reached a hazardous level and traps were raised to a non-sampling position.

Trapping Efficiency

2 trapping efficiency trials have been conducted at Hills Creek Dam. The first being on 2/16/2022 and the second being on 2/25/2022.

During the first Trapping Efficiency trial, a total of 600 juvenile Chinook (parr) were dyed, clipped and released on 02/16/2022 below Hills Creek PWR and 593 below the RO to evaluate the efficiency of the screw trap at those locations. A total of 8 fish were recaptured in the 8ft PWR trap on 2/16/2022 and an additional 4 captured on 2/17/2022 for a total of 12 chinook recaptures in the PWR trap. A total of 19 chinook were captured in the 5ft RO trap on 02/17/2022. No PWR route fish were captured in the RO trap. Route-specific trapping efficiency was 2.0% at the PWR trap and 3.2% at the RO.

Of the fish recaptured,1 was dead and an additional 27 were injured of the total 31 recaptures. Injuries were primarily descaling (16) and fin damage (26). Mt. Hood Environmental staff noted that most fish appeared to have minor descaling with some fin damage, but in good condition overall upon retrieval from the hatchery.

| Hills Creek Dam | Release # | Recapture # | Capture Efficiency |
|-----------------|-----------|-------------|-----------------------|
| PWR Route | 600 | 12 | 2.0% (12/600) |
| RO Trap | 593 | 19 | 3.2% (19/593) |

^{*}Live fish captured at the PWR trap are released just downstream of the PWR trap, upstream of the RO trap and therefore retained in the capture efficiency estimates for the RO Trap if they arrive in the lower trap.

During the second Trapping Efficiency trial, a total of 604 juvenile Chinook (parr) were dyed, clipped and released on 02/25/2022 below Hills Creek PWR and 625 below the RO to evaluate the efficiency of the screw trap at those locations. A total of 5 fish were recaptured in the 8ft PWR trap on 2/26/2022 and an additional 1 captured on 2/27/2022 for a total of 6 chinook recaptures in the PH trap. A total of 6 chinook were captured in the 5ft RO trap on 02/26/2022 and an additional 1 captured on 2/27/2022 for a total of 7 chinook recaptures in the RO trap. No PWR route fish were captured in the RO trap. Route-specific trapping efficiency was 0.99% at the PH trap and 1.12% at the RO.

Of the fish recaptured,1 was dead and an additional 11 were injured of the total 13 recaptures. Injuries were primarily descaling (11) and fin damage (12). Mt. Hood Environmental staff noted that most fish appeared to have minor descaling with some fin damage, but in good condition overall upon retrieval from the hatchery.

^{*}Any dead fish captured at the PWR trap are excluded from the RO trap capture efficiency estimate as they are not alive at time of re-release.

| Hills Creek Dam | Release # | Recapture # | Capture Efficiency |
|-----------------|-----------|-------------|-----------------------|
| PWR Route | 604 | 6 | 0.99% (6/604) |
| RO Trap | 625 | 7 | 1.12% (7/625) |

^{*}Live fish captured at the PWR trap are released just downstream of the PWR trap, upstream of the RO trap and therefore retained in the capture efficiency estimates for the RO Trap if they arrive in the lower trap.

Table 32. Descriptive Statistics of Target Species Captured at Hills Creek Dam Season
To-Date and for the Reporting Period

| | To-Date | | | | | | | | | | | |
|-------------|---------|---------|------------|-----------|--------------|-------|-------|-------------|-------|-------|--|--|
| Site | Route | Species | 125 | Collected | Length (mm)* | | | Weight (g)* | | | | |
| Site | | | Life stage | | Min | Max | Mean | Min | Max | Mean | | |
| Hills Creek | RO | CHS | Parr | 6 | 90.0 | 141.0 | 110.7 | 7.4 | 23.4 | 13.3 | | |
| HIIIS CIEEK | | CHS | Smolt | 62 | 137.0 | 265.0 | 230.9 | 27.35 | 192.3 | 143.9 | | |
| Hills Crook | PWR | CHS | Parr | 7 | 69.0 | 127.0 | 98.1 | 3.7 | 24.5 | 11.2 | | |
| Hills Creek | PWR | CHS | Smolt | 25 | 128.0 | 265.0 | 224.3 | 26.2 | 188.7 | 130.6 | | |

Fish that were missing heads are not included in length and weight calculations. One fish was a head only and could not be assigned a life stage.

| October 1-15, 2022 | | | | | | | | | | | |
|--------------------|-------|---------|------------|-----------|-----|--------------|-------|-----------|-------------|-------|--|
| Cita | Route | Species | Life etama | Collected | Le | Length (mm)* | | | Weight (g)* | | |
| Site | | | Life stage | | Min | Max | Mean | Min | Max | Mean | |
| | | CHS | Parr | 0 | N/A | N/A | N/A | N/A | N/A | N/A | |
| Hills Creek | RO | CHS | Smolt | 1 | 247 | 247 | 247.0 | 174. 0 | 174.0 | 174.0 | |
| Hills Creek | DWD | CHS | Parr | 0 | N/A | N/A | N/A | N/A | N/A | N/A | |
| milis Creek | PWR | CHS | Smolt | 0 | N/A | N/A | N/A | N/A | N/A | N/A | |

Injuries and Copepod Infection

There was 1 Chinook captured in the RO RST. Partial descaling <20% was observed on 0 of 1 Chinook collected at the RO RST (0.0%), and descaling >20% was observed on 1 of the Chinook collected (100.0%). 1 displayed body injuries (100.0%) and 1 had eye injuries (100.0%). 1 of the RO RST Chinook had copepods present in the branchial cavity (100.0%) and 1 had copepods present on fins (100.0%). 0 of the fish captured in the Spill RST displayed Gas Bubble Disease (0.0%).

There were 0 Chinook captured in the Powerhouse channel RST. Partial descaling <20% was observed on 0 of the 0 Chinook collected at the PWR RSTs (0.0%). Descaling >20% was observed on 0 of the

^{*}Any dead fish captured at the PWR trap are excluded from the RO trap capture efficiency estimate as they are not alive at time of re-release.

Chinook collected (0.0%). 0 PWR RST fish had bodily injury (0.0%) and 0 had eye injuries (0.0%). 0 of the fish had copepods present in the branchial cavity (0.0%) and 0 had copepods present on fins (0.0%). 0 fish displayed Gas Bubble Disease (0.0%). There were 0 chinook mortalities collected in the RO RST (0.0%) and 0 in the PWR RST (0.0%). Injuries are displayed in Table 5. To date injury data can be found in Appendix A.

Table 33. Number of Descaled, Bodily/Eye Injured, Copepod Infected and Dead Chinook Salmon for Sampling Period. (Hills Creek Dam)

| Site | Route | # CHS Collected | # DSC* <20% | # DSC* >20% | # with Body Injuries | # with Eye Injuries | # with COP* In B.C. | # with COP* on Fins | Mortalities |
|----------------|-------|--------------------|----------------|----------------|----------------------------|---------------------------|---------------------------|------------------------|-------------|
| Hills Creek | RO | 1 | 0 | 1 | 1 | 1 | 1 | 1 | 0 |
| Hills Creek | PWR | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

^{*}DSC=Descaled, COP=Copepods, B.C.=Branchial Cavity

Collected DNA and Scale Samples

For the reporting period, scales and DNA were collected from 0 Spring Chinook. The other targets captured did not meet length criteria for DNA sampling or were captured before new protocols were put in place.

Non-Target Species

A total of 5 non-target fish were captured at Hills Creek during the reporting period; the data is summarized below in Table 6.

Table 34. Summary of Non-target Species (Hills Creek Dam).

| | | , | 3 - 1 | • | | , |
|--------------------|---------------|-----------------|----------------|------------------|-----------------|------------------------------|
| Species | RO Capture | RO Mortality | PWR Capture | PWR Mortality | Season Total | Season Total Mortality |
| Bluegill | 1 | 1 | 0 | 0 | 55 | 28 |
| Brook Lamprey | 1 | 0 | 0 | 0 | 2 | 0 |
| Bullhead | 0 | 0 | 0 | 0 | 1 | 0 |
| Bull Trout | 0 | 0 | 0 | 0 | 1 | 0 |
| Crappie | 2 | 1 | 0 | 0 | 63 | 41 |
| Longnose Dace | 0 | 0 | 0 | 0 | 2 | 0 |
| Red-Sided Shiner | 1 | 0 | 0 | 0 | 19 | 2 |
| Sculpin | 0 | 0 | 0 | 0 | 46 | 0 |
| Spotted Bass | 0 | 0 | 0 | 0 | 6 | 1 |
| Sucker | 0 | 0 | 0 | 0 | 2 | 1 |
| Mountain Whitefish | 0 | 0 | 0 | 0 | 1 | 1 |
| O. mykiss | 0 | 0 | 0 | 0 | 64 | 22 |
| Unknown | 0 | 0 | 0 | 0 | 4 | 2 |
| Totals | 5 | 2 | 0 | 0 | 287 | 100 |

Stream Statistics

Basic stream statistics at the Hills Creek site were calculated from data downloaded from the U.S. Geological Survey stream gauge numbers 14145110 and 14145500. Gauge height (feet) is the only metric provided at this gauge. Total dissolved gas saturation data was received from gauge 14145500, 1.4 rkms downstream of the trap. During the reporting period, daily maximum values for instantaneous gauge height ranged from 1,224.3 feet to 1,224.6 feet (mean: 1,224.6 feet). Figure 50 shows instantaneous gauge height.

Total dissolved gas saturation ranged from 101 to 105% (mean: 102.8%) during the reporting period. Figure 51 shows total dissolved gas saturation.

Stream temperatures were recorded every 2 hours for the both the RO RST and the PWR RST (Figures 52 and 53). The probes operated normally throughout the reporting period.

Flows through the PWR and RO during the reporting period averaged 0 and 771.4 cubic feet per second (cfs) respectively (Figure 54). Catch per unit of effort (CPUE) data are summarized in Table 7. Discharge and capture data for the duration of monitoring efforts at this location are provided in Appendix B.

Table 35. Summary of Chinook CPUE, Hills Creek Dam.

| | Chinook | | | | | | | | | |
|----------------|----------|-----------|--|--|--|--|--|--|--|--|
| Description | RO (5ft) | PWR (8ft) | | | | | | | | |
| Catch | 1 | 0 | | | | | | | | |
| Effort (hrs) | 239.3 | 239.2 | | | | | | | | |
| CPUE (fish/hr) | 0.004 | 0 | | | | | | | | |




Figure 50. Gauge Height (feet); below Hills Creek Dam PWR - Middle Fork Willamette River




Figure 51. Total Dissolved Gas Saturation (%); below Hills Creek Dam – Middle Fork Willamette River




Figure 52. Temperature at RO RST (Hills Creek Dam)




Figure 53. Temperature at Powerhouse RST (Hills Creek Dam)




Figure 54. Hourly Flows PWR vs. RO (Hills Creek Dam)

Issues Encountered

The Cedar Creek Fire is still active Southeast of the town of Oakridge near two of our sites on the Middle Fork Willamette River. Air quality exceeded 500 ppm this reporting period. Hills Creek Dam, Lookout Point Head of Reservoir, Lookout Dam Tailrace, and Dexter Dam Tailrace traps were raised to a non-sampling position on October 10th. These traps were not returned to a sampling position for the remainder of the reporting period as AQI remained in the hazardous category. The flow in the Cougar Dam powerhouse channel was lowered on 10/8/2022 in order to prioritize flow from the regulating outlet. The powerhouse traps were raised on 10/8/2022 in anticipation of this event to avoid damage to the cones. The flow is too low to continue fishing the Cougar Dam powerhouse channel.

Upcoming USACE Support Services

None at this time.

Appendix A

Chinook (CHS)

| | | | | | | Ch | ino | ok Ini | uries | to- | date | | | | | | | | | | | | |
|----------------------|--------------|------|------|-----|-----|-----|-----|--------|-------|-----|------|-----|----|---|-----|-----|-----|-----|-----|----|-----|-----|-----|
| Site/Trap/Life Stage | e Total Fish | MUNK | DS<2 | BLO | EYB | FUN | BKD | COP | DS>2 | PRD | HD | НВО | ВО | 오 | BVT | НВР | BRU | TEA | OPD | Z | FVB | POP | GBD |
| Big Cliff Dam | 1107 | | 683 | 4 | 75 | 3 | 1 | 846 | 253 | 4 | 555 | 5 | 10 | 4 | 24 | 6 | 66 | 36 | 118 | 69 | 63 | 13 | 32 |
| 8 ft | 1107 | | 683 | 4 | 75 | 3 | 1 | 846 | 253 | 4 | 555 | 5 | 10 | 4 | 24 | 6 | 66 | 36 | 118 | 69 | 63 | 13 | 32 |
| Adult | 1 | | 1 | | | | | | | | 1 | | | | | | 1 | 1 | 1 | | 1 | | |
| Parr | 24 | | 6 | | 1 | 1 | | 15 | 3 | | 6 | | | | | 2 | | | 1 | | | | |
| Smolt | 1071 | | 676 | 4 | 73 | 2 | | 831 | 249 | 4 | 547 | 5 | 10 | 2 | 24 | 4 | 64 | 35 | 115 | 68 | 62 | 12 | 32 |
| Unknown | 2 | | | | | | | | | | | | | 2 | | | | | | | | 1 | |
| Fry | 9 | | | | 1 | | 1 | | 1 | | 1 | | | | | | 1 | | 1 | 1 | | | |
| Foster Dam HOR | | | | | | | | | | | | | | | | | | | | | | | |
| 5 ft | 66 | | 9 | | 1 | | | | | | 2 | | | | | | | | | | | | |
| Parr | 8 | | 5 | | | | | | | | 1 | | | | | | | | | | | | |
| Smolt | 3 | | 3 | | | | | | | | 1 | | | | | | | | | | | | |
| Fry | 55 | | 1 | | 1 | | | | | | | | | | | | | | | | | | |
| Unknown | 2 | | | | | | | | | | | | | 2 | | | | | | | | | |
| Cougar Dam | | | | | | | | | 209 | | | | | | | | | | | | | | |
| RO | 431 | | 251 | 3 | 64 | 2 | 3 | 299 | 89 | 3 | 186 | 2 | 1 | | 8 | 2 | 20 | 9 | 48 | 27 | 13 | 4 | 30 |
| Parr | 135 | | 70 | | 23 | | | 77 | 25 | | 48 | | | | 1 | | 5 | 2 | 11 | 7 | | | 6 |
| Smolt | 275 | | 181 | 3 | 38 | 2 | 3 | 222 | 63 | 3 | 137 | 2 | 1 | | 7 | 2 | 14 | 7 | 36 | 18 | 13 | 4 | 24 |
| Fry | 21 | | | | 3 | | | | 1 | | 1 | | | | | | 1 | | 1 | 2 | | | |
| PH | 1188 | 10 | 513 | 5 | 32 | 3 | | 568 | 120 | | 306 | | 5 | 2 | 34 | 2 | 27 | 29 | 49 | 21 | 62 | 4 | 4 |
| Parr | 266 | | 146 | | 11 | 1 | | 104 | 25 | | 67 | | 1 | | 5 | | 3 | 5 | 11 | 6 | 5 | | |
| Smolt | 538 | | 365 | 5 | 17 | 2 | | 464 | 89 | | 234 | | 4 | | 28 | 2 | 22 | 17 | 35 | 12 | 57 | 1 | 4 |
| Unknown | 2 | | | | | | | | | | | | | 2 | | | | | | | | | |
| Fry | 382 | 10 | 2 | | 4 | | | | 6 | | 5 | | | | 1 | | 2 | 7 | 3 | 3 | | 3 | |
| Cougar Dam HOR | | | | | | | | | | | | | | | | | | | | | | | |
| 5 ft | 632 | 3 | 50 | | | | | 6 | 1 | 1 | 30 | | | | | | | 4 | 7 | 6 | 1 | 1 | |
| Parr | 100 | | 47 | | | | | 6 | 1 | | 22 | | | | | | | 4 | | | 1 | | |
| Smolt | 4 | | 1 | | | | | | | | | | | | | | | | | | | | |
| Fry | 528 | 3 | 2 | | | | | | | 1 | 8 | | | | | | | | 7 | 6 | | 1 | |
| Fall Creek HOR | | | | | | | | | | | | | | | | | | | | | | | |
| 8 ft | 7 | | 3 | | | | | 2 | | | 1 | | | | | | | | | | | | |
| Parr | 2 | | 2 | | | | | 1 | | | 1 | | | | | | | | | | | | |
| Smolt | 5 | | 1 | | | | | 1 | | | | | | | | | | | | | | | |
| Dexter Dam Tail. | 98 | | | | | | | | | | 46 | | | | | | | | | | | | |
| 5 ft | 98 | | 59 | | 6 | | | 11 | 23 | | 46 | | | | 1 | | 3 | 4 | 6 | 6 | 4 | | 20 |
| Parr | 18 | | 6 | | 3 | | | 2 | 6 | | 8 | | | | | | | 2 | 3 | 2 | | | 6 |
| Smolt | 77 | | 53 | | 3 | | | 9 | 17 | | 37 | | | | 1 | | 3 | 2 | 3 | 4 | 4 | | 14 |
| Fry | 3 | | | | | | | | | | 1 | | | | | | | | | | | | |

Chinook (CHS)

| | | | | Chino | ok Inj | uries | to-date | | | | | | | | | | | |
|----------------------|-----------------------------|------|------------|------------|--------|-------|------------|-----|-----|-----|-----|-----|-----|-----|---|-----|-----|-----|
| Site/Trap/Life Stage | Y Z ⊃ Total Fish ∑ | DS<2 | BLO EYB | FUN BKD | COP | DS>2 | PRD FID | HBO | 위 | BVT | НВР | BRU | TEA | OPD | Z | FVB | POP | GBD |
| Lookout Dam Tail. | 78 | 41 | 1 15 | 1 | 14 | 28 | 44 | | | 2 | 2 | 9 | 1 | 15 | 9 | 6 | | 8 |
| PH1 | 28 | 11 | 1 7 | | 7 | 14 | 19 | | | 2 | 2 | 6 | | 8 | 2 | 1 | | 2 |
| Parr | 3 | | 1 | | | 2 | 2 | | | | 1 | 1 | | 1 | | | | |
| Smolt | 25 | 11 | 1 6 | | 7 | 12 | 17 | | | 2 | 1 | 5 | | 7 | 2 | 1 | | 2 |
| PH 2 | 12 | 10 | 3 | | 2 | 2 | 9 | | | | | 1 | 1 | 2 | 4 | 1 | | |
| Parr | 4 | 4 | 3 | | | 1 | 4 | | | | | 1 | 1 | 1 | 3 | | | |
| Smolt | 8 | 6 | | | 2 | 1 | 5 | | | | | | | 1 | 1 | 1 | | |
| Spill | 38 | 20 | 5 | 1 | 5 | 12 | 16 | | | | | 2 | | 5 | 3 | 4 | | 6 |
| Parr | 6 | 1 | | | | 3 | 1 | | | | | | | | | | | |
| Smolt | 32 | 19 | 5 | 1 | 5 | 9 | 15 | | | | | 2 | | 5 | 3 | 4 | | 6 |
| Lookout Point HOR | 103 | 31 | | | 2 | | 12 | | | | | 1 | | 1 | | | | |
| 5 ft | 103 | 31 | | | 2 | | 12 | | | | | 1 | | 1 | | | | |
| Parr | 59 | 29 | | | 1 | | 9 | | | | | 1 | | 1 | | | | |
| Smolt | 3 | 1 | | | 1 | | 1 | | | | | | | | | | | |
| Fry | 41 | 1 | | | | | 2 | | | | | | | | | | | |
| Hills Creek Dam | 101 | 1 55 | | | | 36 | | | 8 1 | | | | | | | | | 2 |
| RO | 69 | 1 37 | 6 | | 55 | 25 | 18 | | 6 | 22 | 6 | 2 | 2 | 6 | 2 | 2 | | 2 |
| Parr | 6 | 1 | | | 1 | | | | | | | 1 | | | | | | |
| Smolt | 63 | 1 36 | 6 | | 54 | 25 | 18 | | 6 | 22 | 6 | 1 | 2 | 6 | 2 | 2 | | 2 |
| PH | 32 | 18 | 6 | | 21 | 11 | 7 | | 2 1 | . 6 | 2 | 1 | 1 | 1 | 2 | 3 | | |
| Parr | 7 | 4 | | | 1 | 1 | | | | | 1 | | | | 1 | | | |
| Smolt | 25 | 14 | 6 | | 20 | 10 | 7 | | 2 1 | 6 | 1 | 1 | 1 | 1 | 1 | 3 | | |


Steelhead (O. mykiss)


| | | | <u> </u> | emeau | (O. III) | (KISS) | | | | | | |
|----------------------|--------------|------|------------|------------|-------------|------------|----------|------------|------------|-----|------------|-----|
| | | | 0. r | nykiss Inj | uries To | Date | | | | | | |
| Site/Trap/Life Stage | Total Fish ≥ | DS<2 | EYB FUN | ВКР | DS>2 PRD | FID HBO | BO HO | BVT HBP | BRU TEA | OPD | HIN FVB | POP |
| Big Cliff Dam | 62 | | | 10 | | | | | 4 3 | | | 1 |
| 8 ft | 62 | 17 | 1 | 10 | 5 | 15 | 1 | 1 | 4 3 | 5 | 6 | 1 |
| Adult | 1 | | | | 1 | 1 | 1 | | | | | |
| Parr | 17 | 4 | 1 | | | 4 | | | | | 1 | |
| Smolt | 16 | 10 | | 10 | 4 | 9 | | 1 | 3 3 | 5 | 5 | 1 |
| Fry | 28 | 3 | | | | 1 | | | 1 | | | |
| Green Peter Tail. | | | | | | | | | | | | 5 |
| 8 ft | 6 | 3 | 1 | 1 | 2 | 4 | | | 4 | 1 | 1 | 5 |
| Smolt | 6 | 3 | 1 | 1 | 2 | 4 | | | 4 | 1 | 1 | 5 |
| Foster Dam HOR | 90 | 25 | | | | 11 | | | 1 | | | |
| 5 ft | 90 | 25 | | 1 | | 11 | | | 1 | | | |
| Adult | 7 | 1 | | | | 2 | | | | | | |
| Parr | 17 | 7 | | 1 | | 2 | | | | | | |
| Smolt | 35 | 17 | | | | 7 | | | 1 | | | |
| Fry | 31 | | | | | | | | | | | |

Steelhead (O. mykiss)

| | Steemeau (O. mykiss) | | | | | | | | | | | | | | | | | | | | | | |
|--------------------|--|------|------|-----|-----|-----|-----|-----|------|-----|-----|-----|----|---|-----|-----|-----|-----|-----|--------|-----|-----|-----|
| | O. mykiss Injuries During Reporting Period (10-1-2022 to 10-15-2022) | | | | | | | | | | | | | | | | | | | | | | |
| Site/Trap/Life Sta | ige Total Fish | MUNK | DS<2 | BLO | EYB | FUN | BKD | COP | DS>2 | PRD | FID | НВО | BO | 오 | BVT | НВР | BRU | TEA | OPD | Z I | FVB | POP | GBD |
| Big Cliff Dam | | | | | | | | | | | | | | | | | | | | | | | |
| 8 ft | 2 | | | | | | | | | | | | | | | | | | | | | | |
| Parr | 1 | | | | | | | | | | | | | | | | | | | | | | |
| Fry | 1 | | | | | | | | | | | | | | | | | | | | | | |


Appendix B







Green Peter Dam Operational and Green Peter Tailrace- Middle Santiam River Capture Data Since





















South Fork McKenzie above Cougar Dam Discharge and Cougar Dam Head of Reservoir Capture Data Since Start of Monitoring

















Note: Gauge height data was taken from USGS stream gauge number 14150290, 1.2 rkms downstream.




















Appendix C

Hills Creek Trapping Efficiency 1/6/2022

| Hills Creek Dam | Release # | Recapture # | Capture Efficiency |
|-----------------|------------------|-------------|-----------------------|
| PH Route | 596 | 20 | 3.36% (20/596) |
| DO Tran | RO Route- 605 | 13 | 2.15% (13/605) |
| RO Trap | PH Route- 592 | 5 | 0.84% (5/592) |

^{*}Live fish captured at the PH trap are released just downstream of the PH trap, upstream of the RO trap and therefore retained in the capture efficiency estimates for the RO Trap if they arrive in the lower trap.

*Any dead fish captured at the PH trap are excluded from the RO trap capture efficiency estimate as they are not alive at time of re-release.

| Cougar Dam | Release # | Recapture # | Capture Efficiency |
|------------|-----------|-------------|--------------------|
| PH Route | 405 | 40 | 9.88% (40/405) |
| RO Route | 410 | 28 | 6.83% (28/410) |

| Dexter Dam | Release # | Recapture # | Capture Efficiency |
|------------|-----------|-------------|-----------------------|
| Spill | 988 | 2 | 0.2% (2/988) |
| Powerhouse | N/A | N/A | N/A |

| Green Peter Dam Tailrace- Middle Santiam | Release # | Recapture # | Capture Efficiency |
|--|-----------|-------------|-----------------------|
| 8ft Trap | 643 | 4 | 0.62% (4/643) |

| Big Cliff Dam | Release # | Recapture # | Capture Efficiency |
|---------------|-----------|-------------|-----------------------|
| 8ft Trap | 996 | 40 | 4.01% (40/996) |

| Cougar Dam Head of Reservoir | Release # | Recapture # | Capture Efficiency |
|------------------------------|-----------|-------------|-----------------------|
| 5ft trap | 806 | 41 | 5.1% (41/806) |

| Cougar Dam Head of Reservoir | Release # | Recapture # | Capture Efficiency |
|------------------------------|-----------|-------------|-----------------------|
| 5ft trap | 515 | 7 | 1.4% (7/515) |

| Cougar Dam | Release # | Recapture # | Capture Efficiency |
|------------|-----------|-------------|-----------------------|
| PH Route | 357 | 62 | 17.37% (62/357) |
| RO Route | 378 | 21 | 5.56% (21/378) |

| Dexter Dam | Release # | Recapture # | Capture Efficiency |
|------------|-----------|-------------|-----------------------|
| Spill | 1000 | 43 | 4.3% (43/1000) |
| Powerhouse | N/A | N/A | N/A |

| Lookout Point Head of Reservoir | Release # | Recapture # | Capture Efficiency |
|------------------------------------|-----------|-------------|-----------------------|
| 04/05/2022 | 993 | 53 | 5.3% (53/993) |
| 04/14/2022 | 989 | 19 | 1.9% (19/989) |

| Fall Creek Dam | Release # | Recapture # | Capture Efficiency |
|----------------|-----------|-------------|-----------------------|
| RO | 518 | 11 | 2.1% (11/518) |

| Fall Creek Dam | Release # | Recapture # | Capture Efficiency |
|----------------|-----------|-------------|-----------------------|
| RO | 513 | 0 | 0% (0/513) |

| Cougar Dam | Release # | Recapture # | Capture Efficiency |
|------------|-----------|-------------|-----------------------|
| RO Route | 993 | 63 | 6.34% (63/993) |

| Dexter Dam | Release # | Recapture # | Capture Efficiency |
|------------|-----------|-------------|-----------------------|
| Spill | 1019 | 67 | 6.6% (67/1,019) |
| Powerhouse | N/A | N/A | N/A |

| Lookout Point Head of Reservoir | Release # | Recapture # | Capture Efficiency |
|------------------------------------|-----------|-------------|-----------------------|
| 05/24/2022 | 1007 | 125 | 12.4% (125/1007) |

| Big Cliff Dam | Release # | Recapture # | Capture Efficiency |
|---------------|-----------|-------------|-----------------------|
| 8ft Trap | 1000 | 21 | 2.1% (21/1000) |

| Cougar Dam | Release # | Recapture # | Capture Efficiency |
|------------|-----------|-------------|-----------------------|
| PH Route | 500 | 148 | 29.6% (148/500) |

| Cougar Dam Head of Reservoir | Release # | Recapture # | Capture Efficiency |
|------------------------------|-----------|-------------|-----------------------|
| 5ft trap | 551 | 56 | 10.2% (56/551) |

| Big Cliff Dam | Release # | Recapture # | Capture Efficiency |
|---------------|-----------|-------------|-----------------------|
| 8ft Trap | 1000 | 92 | 9.2% (92/1000) |

| Cougar Dam | Release # | Recapture # | Capture Efficiency |
|------------|-----------|-------------|-----------------------|
| PH Route | 501 | 31 | 6.2% (31/501) |

Appendix D

Summary of Project PIT Tagged Fish for Reporting Period

| Site | Trap | # of PIT Tagged Fish |
|---|-------|----------------------|
| Big Cliff Dam | 8 ft | 0 |
| Foster Dam Head of Reservoir- South Santiam | 5 ft | 0 |
| Cougar Dam | PWR | 85 |
| Cougar Dam | RO | 5 |
| Cougar Dam Head of Reservoir | 5 ft | 10 |
| Green Peter Tailrace- Middle Santiam | 8 ft | 0 |
| Dexter Dam Tailrace | 5 ft | 0 |
| Lookout Point Head of Reservoir | 5 ft | 0 |
| Lookout Dam Tailrace | Spill | 0 |
| Lookout Dam Tailrace | PWR | 0 |
| Hills Creek Dam Tailrace | PWR | 0 |

| Hills Creek Dam Tailrace | RO | 1 |
|--------------------------|----|---|
| | | |

To Date Summary of Captured Fish Containing PIT Tags

| Site | Trap | PIT Tag # | Date | Species |
|------------------------------|------|----------------|-----------|-----------|
| Cougar Dam | RO | 3DD.0077780789 | 1/8/2022 | Chinook |
| Cougar Dam | RO | 384.36F2B2C55F | 1/14/2022 | Chinook |
| Cougar Dam | PH | 3DD.003DA4DC74 | 3/3/2022 | Chinook |
| Cougar Dam | PH | 3DD.003E14CA70 | 3/4/2022 | Chinook |
| Cougar Dam | PH | 384.36F2B2C5D2 | 3/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003E14CC20 | 3/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003E14C9D6 | 3/6/2022 | Chinook |
| Cougar Dam | PH | 3DD.003E14CD8D | 3/8/2022 | Chinook |
| Cougar Dam | RO | 3DD.003BD59645 | 4/7/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BD21883 | 7/23/2022 | O. mykiss |
| Cougar Dam | PH | 3DD.003BD21883 | 7/26/2022 | O. mykiss |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE1849 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE22CE | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2AAF | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE1885 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE24AD | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2293 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE26D4 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2422 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2AB1 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE227B | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE22B9 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE24D6 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE22B8 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE223A | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE18C0 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE1965 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE224D | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE242B | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2464 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE244B | 9/23/2022 | Chinook |

| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2443 | 9/23/2022 | Chinook |
|------------------------------|------|----------------|-----------|---------|
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE26F9 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2449 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2519 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2517 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2AF3 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE18D2 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE18E3 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE185B | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE223F | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE270D | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE16F5 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2284 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE175F | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2252 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE223B | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE240E | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE26D0 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2253 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2489 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE244E | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE190D | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE18D8 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2AC4 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE176A | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE1917 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2AEF | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE192E | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2266 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE1916 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE16D7 | 9/23/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2216 | 9/24/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE1741 | 9/24/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2AFB | 9/24/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE16BC | 9/25/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE26CC | 9/27/2022 | Chinook |

| Cougar Dam | PH | 3DD.003BEE22C0 | 10/4/2022 | Chinook |
|------------------------------|------|----------------|-----------|---------|
| Cougar Dam | PH | 3DD.003BEE2C55 | 10/7/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE192F | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE241F | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2711 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE26FE | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE18AA | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE22C8 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE1933 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE224B | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2474 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE170A | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE16F8 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2291 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE1953 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE226B | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE1757 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2506 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE225E | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE18D4 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE1857 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2418 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE18B6 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2492 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE18EA | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE26BE | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE172B | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2713 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE26C4 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE1706 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE18CA | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2AA7 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE18E2 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE18F0 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE26EA | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE176B | 10/6/2022 | Chinook |

| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE1957 | 10/6/2022 | Chinook |
|------------------------------|------|----------------|------------|---------|
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE226D | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE1901 | 10/6/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE18B5 | 10/7/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE16D3 | 10/7/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE16EC | 10/7/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE186E | 10/7/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE23FC | 10/7/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE1910 | 10/8/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE192B | 10/9/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE16C5 | 10/10/2022 | Chinook |

List of EAS PIT Tagged Fish for Reporting Period

| Site | Trap | PIT Tag # | Date | Species |
|------------|------|----------------|-----------|---------|
| Cougar Dam | PH | 3DD.003BEE2C35 | 10/3/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2C8F | 10/3/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2C52 | 10/3/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2C41 | 10/3/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE264E | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2647 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE25FF | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE263D | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2642 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2646 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2616 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE25F2 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2610 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE263C | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE25F3 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE262C | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2615 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2635 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE263F | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2629 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2636 | 10/4/2022 | Chinook |

| Cougar Dam | PH | 3DD.003BEE260D | 10/4/2022 | Chinook |
|------------|----|----------------|-----------|---------|
| Cougar Dam | PH | 3DD.003BEE264B | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE262B | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2648 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2652 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE25FC | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2603 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2623 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE260F | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2604 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE25FB | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2608 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2602 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE260B | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE260E | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A37 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A0C | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE29E5 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A18 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A15 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE29E2 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A0F | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE29FC | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A03 | 10/4/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A2B | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A00 | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE29E9 | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE29FD | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A25 | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A17 | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE29F7 | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A0D | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A14 | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE29E1 | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A2D | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A32 | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE29F5 | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A0E | 10/5/2022 | Chinook |

| Cougar Dam | PH | 3DD.003BEE2A27 | 10/5/2022 | Chinook |
|------------------------------|------|----------------|------------|---------|
| Cougar Dam | PH | 3DD.003BEE2A24 | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A20 | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A1E | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A23 | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A2A | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2A1D | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE29EE | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE29F8 | 10/5/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE1682 | 10/6/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE1654 | 10/6/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE168C | 10/6/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE1687 | 10/6/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE1697 | 10/6/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE1683 | 10/6/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE165E | 10/6/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE1695 | 10/6/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE1663 | 10/6/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE1669 | 10/6/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE166B | 10/6/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE1652 | 10/6/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE1666 | 10/6/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2C85 | 10/7/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2C3E | 10/7/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2C68 | 10/7/2022 | Chinook |
| Cougar Dam | PH | 3DD.003BEE2C79 | 10/7/2022 | Chinook |
| Cougar Dam | RO | 3DD.003BEE169A | 10/8/2022 | Chinook |
| Cougar Dam | RO | 3DD.003BEE169F | 10/8/2022 | Chinook |
| Cougar Dam | RO | 3DD.003BEE169D | 10/8/2022 | Chinook |
| Cougar Dam | RO | 3DD.003BEE2C2F | 10/10/2022 | Chinook |
| Cougar Dam | RO | 3DD.003BEE2C46 | 10/10/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE25FD | 10/1/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2C90 | 10/2/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2C66 | 10/3/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2C6D | 10/3/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2C7A | 10/3/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2C4E | 10/3/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2A11 | 10/4/2022 | Chinook |

| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE29DA | 10/5/2022 | Chinook |
|------------------------------|------|----------------|------------|---------|
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE29F3 | 10/5/2022 | Chinook |
| Cougar Dam Head of Reservoir | 5 ft | 3DD.003BEE2A36 | 10/5/2022 | Chinook |
| Hills Creek Dam | RO | 3DD.003BEE1CC1 | 10/10/2022 | Chinook |